=> d stat que L52

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS is strictly prohibited.

FILE COVERS 1907 - 2 Jan 2009 VOL 150 ISS 2 FILE LAST UPDATED: 1 Jan 2009 (20090101/ED)

ZCAplus now includes complete International Patent Classification (IPC) reclassification data for the third quarter of 2008.

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file contains CAS Registry Numbers for easy and accurate substance identification.

'OBI' IS DEFAULT SEARCH FIELD FOR 'ZCAPLUS' FILE

L44	22	SEA	FILE=ZCAPLUS	SPE=ON	ABB=ON	PLU=ON	RASNETSOV L?/AU
L45	55	SEA	FILE=ZCAPLUS	SPE=ON	ABB=ON	PLU=ON	SHVARTSMAN I?/AU
L46	13	SEA	FILE=ZCAPLUS	SPE=ON	ABB=ON	PLU=ON	LYALINA I?/AU
L47	19	SEA	FILE=ZCAPLUS	SPE=ON	ABB=ON	PLU=ON	RASNETSOVA B?/AU
L49	15	SEA	FILE=ZCAPLUS	SPE=ON	ABB=ON	PLU=ON	L44 AND (L45 OR L46
		OR I	L47)				
L50	1	SEA	FILE=ZCAPLUS	SPE=ON	ABB=ON	PLU=ON	L45 AND (L46 OR L47)
L51	2	SEA	FILE=ZCAPLUS	SPE=ON	ABB=ON	PLU=ON	L46 AND L47
L52	15	SEA	FILE=ZCAPLUS	SPE=ON	ABB=ON	PLU=ON	L49 OR L50 OR L51
=> d stat	que L	55					
			FILE=ZCAPLUS		ABB=ON	PLU=ON	
			FILE=ZCAPLUS		ABB=ON	PLU=ON	
L46			FILE=ZCAPLUS		ABB=ON	PLU=ON	
L47			FILE=ZCAPLUS		ABB=ON	PLU=ON	
			FILE=ZCAPLUS				?FULLEREN?/BI
L55	11		FILE=ZCAPLUS	SPE=ON	ABB=ON	PLU=ON	(L44 OR L45 OR L46 OR
		L47) AND L54				
		_					
=> d stat							
			FILE=ZCAPLUS		ABB=ON	PLU=ON	
L44			FILE=ZCAPLUS		ABB=ON	PLU=ON	
L45			FILE=ZCAPLUS		ABB=ON	PLU=ON	
L46			FILE=ZCAPLUS		ABB=ON	PLU=ON	
L47			FILE=ZCAPLUS		ABB=ON	PLU=ON	
L49	15		FILE=ZCAPLUS	SPE=ON	ABB=ON	PLU=ON	L44 AND (L45 OR L46
			L47)				
L50			FILE=ZCAPLUS		ABB=ON	PLU=ON	
L51			FILE=ZCAPLUS		ABB=ON	PLU=ON	
L52			FILE=ZCAPLUS		ABB=ON	PLU=ON	
			FILE=ZCAPLUS		ABB=ON	PLU=ON	
L55			FILE=ZCAPLUS	SPE=ON	ABB=ON	PLU=ON	(L44 OR L45 OR L46 OR
) AND L54				
L57	2	SEA	FILE=ZCAPLUS	SPE=ON	ABB=ON	PLU=ON	L5 AND (L52 OR L55)

```
=> s L52 or L55 or L52
           18 L52 OR L55 OR L52
=> file medline embase biosis wpix
FILE 'MEDLINE' ENTERED AT 16:12:21 ON 02 JAN 2009
FILE 'EMBASE' ENTERED AT 16:12:21 ON 02 JAN 2009
Copyright (c) 2009 Elsevier B.V. All rights reserved.
FILE 'BIOSIS' ENTERED AT 16:12:21 ON 02 JAN 2009
Copyright (c) 2009 The Thomson Corporation
FILE 'WPIX' ENTERED AT 16:12:21 ON 02 JAN 2009
COPYRIGHT (C) 2009 THOMSON REUTERS
=> d stat que L53
L44
           22 SEA FILE=ZCAPLUS SPE=ON ABB=ON PLU=ON RASNETSOV L?/AU
L45
            55 SEA FILE-ZCAPLUS SPE-ON ABB-ON PLU-ON SHVARTSMAN I?/AU
           13 SEA FILE=ZCAPLUS SPE=ON ABB=ON PLU=ON LYALINA I?/AU
L46
L47
           19 SEA FILE=ZCAPLUS SPE=ON ABB=ON PLU=ON RASNETSOVA B?/AU
L49
           15 SEA FILE=ZCAPLUS SPE=ON ABB=ON PLU=ON L44 AND (L45 OR L46
             OR L47)
L50
            1 SEA FILE=ZCAPLUS SPE=ON ABB=ON PLU=ON L45 AND (L46 OR L47)
L51
            2 SEA FILE=ZCAPLUS SPE=ON ABB=ON PLU=ON L46 AND L47
           15 SEA FILE=ZCAPLUS SPE=ON ABB=ON PLU=ON L49 OR L50 OR L51
L52
           14 SEA L52
L53
=> d stat que L58
L44 22 SEA FILE=ZCAPLUS SPE=ON ABB=ON PLU=ON RASNETSOV L?/AU
L45
            55 SEA FILE=ZCAPLUS SPE=ON ABB=ON PLU=ON SHVARTSMAN I?/AU
           13 SEA FILE=ZCAPLUS SPE=ON ABB=ON PLU=ON LYALINA I?/AU
L46
           19 SEA FILE=ZCAPLUS SPE=ON ABB=ON PLU=ON RASNETSOVA B?/AU
L47
L58
            9 SEA (L44 OR L45 OR L46 OR L47) AND ?FULLEREN?
=> d stat que L59
L44 22 SEA FILE=ZCAPLUS SPE=ON ABB=ON PLU=ON RASNETSOV L?/AU
           55 SEA FILE=ZCAPLUS SPE=ON ABB=ON PLU=ON SHVARTSMAN 1?/AU
13 SEA FILE=ZCAPLUS SPE=ON ABB=ON PLU=ON LYALINA 1?/AU
L45
L46
L47
           19 SEA FILE=ZCAPLUS SPE=ON ABB=ON PLU=ON RASNETSOVA B?/AU
L49
           15 SEA FILE=ZCAPLUS SPE=ON ABB=ON PLU=ON L44 AND (L45 OR L46
              OR L47)
            1 SEA FILE=ZCAPLUS SPE=ON ABB=ON PLU=ON L45 AND (L46 OR L47)
L51
            2 SEA FILE-ZCAPLUS SPE-ON ABB-ON PLU-ON L46 AND L47
L52
            15 SEA FILE=ZCAPLUS SPE=ON ABB=ON PLU=ON L49 OR L50 OR L51
           14 SEA L52
1.53
L58
            9 SEA (L44 OR L45 OR L46 OR L47) AND ?FULLEREN?
L59
            3 SEA (L53 OR L58) AND AMINO ACID?
=> s L53 or L58 or L59
L61 14 L53 OR L58 OR L59
=> dup rem L60 L61
FILE 'ZCAPLUS' ENTERED AT 16:12:55 ON 02 JAN 2009
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
```

COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

```
FILE 'WPIX' ENTERED AT 16:12:55 ON 02 JAN 2009
COPYRIGHT (C) 2009 THOMSON REUTERS
PROCESSING COMPLETED FOR L60
```

PROCESSING COMPLETED FOR L61

L62 19 DUP REM L60 L61 (13 DUPLICATES REMOVED)

ANSWERS '1-18' FROM FILE ZCAPLUS

ANSWER '19' FROM FILE WPIX

=> d ibib abs hitind L62 1-18; d iall hit L62 19

L62 ANSWER 1 OF 19 ZCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 1

ACCESSION NUMBER: 2008:1359611 ZCAPLUS Full-text

DOCUMENT NUMBER: 149:519138

TITLE: Ophthalmological gel and a method for the use thereof INVENTOR(S): Rasnetsov. Lev Davidovich: Shvartsman, Takov

Yudelevich; Yashnova, Olga Konstantinovna; Melnikova,

Nina Borisovna; Kolchik, Olga Vladimirovna; Gusikhina, Maria Sergeevna

PATENT ASSIGNEE(S): Russia

SOURCE: PCT Int. Appl., 25pp.

CODEN: PIXXD2
DOCUMENT TYPE: Patent

DOCUMENT TYPE:

LANGUAGE: Russian

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.			KIN	KIND DATE			APPLICATION NO.					DATE					
		A1 20081113			WO 2008-RU259					20080423							
	W:	ΑE,	AG,	AL,	AM,	AO,	AT,	AU,	AZ,	BA,	BB,	BG,	BH,	BR,	BW,	BY,	BZ,
		CA,	CH,	CN,	CO,	CR,	CU,	CZ,	DE,	DK,	DM,	DO,	DZ,	EC,	EE,	EG,	ES,
		FI,	GB,	GD,	GE,	GH,	GM,	GT,	HN,	HR,	HU,	ID,	IL,	IN,	IS,	JP,	KE,
		KG,	KM,	KN,	KΡ,	KR,	KZ,	LA,	LC,	LK,	LR,	LS,	LT,	LU,	LY,	MA,	MD,
		ME,	MG,	MK,	MN,	MW,	MX,	MY,	ΜZ,	NA,	NG,	NI,	NO,	NZ,	OM,	PG,	PH,
		PL,	PT,	RO,	RS,	SC,	SD,	SE,	SG,	SK,	SL,	SM,	SV,	SY,	ΤJ,	TM,	TN,
		TR,	TT,	TZ,	UA,	UG,	US,	UZ,	VC,	VN,	ZA,	ZM,	ZW				
	RW:	AT,	BE,	BG,	CH,	CY,	CZ,	DE,	DK,	EE,	ES,	FI,	FR,	GB,	GR,	HR,	HU,
		IE,	IS,	IT,	LT,	LU,	LV,	MC,	MT,	NL,	NO,	PL,	PT,	RO,	SE,	SI,	SK,
		TR,	BF,	ВJ,	CF,	CG,	CI,	CM,	GA,	GN,	GQ,	GW,	ML,	MR,	NE,	SN,	TD,
		TG,	BW,	GH,	GM,	KE,	LS,	MW,	MZ,	NA,	SD,	SL,	SZ,	TZ,	UG,	ZM,	ZW,
		AM,	ΑZ,	BY,	KG,	KZ,	MD,	RU,	ΤJ,	TM							
RU	2340	327			C1		2008	1210		RU 2	007-	1167	79		2	0070	503

PRIORITY APPLN. INFO::

RU 2007-116779 A 20070503

AB The invention relates to medicine. The inventive ophthalmol, gel comprises

0.2-0.5% low-cross-linked polyacrylic acid and/ or derivs, thereof, preservatives, stabilizers, a medicinal substance selected from a reparant group and clean water, and has a pH value within the range of lachrymal liquid The gel contains $1-(\beta-\text{csyethyl})-4,6-\text{dimethyl-1},2-\text{dihydro-2-oxypyrimidine}$ (xymedon) in the form of a medicinal substance. The inventive gel production method consists in adding an aqueous 10-30% polyethylene oxide solution into a dry powder of low-cross-linked polyacrylic acid associated with rapid agitation, polyethylene oxide being taken at least in a tenfold excess with respect to the mass of the powder, in adding, while agitating, clean water in a quantity equal to 70-90% the total mass of the gel, in adjusting a pH value to a value of 6.0-7.0 by means of a sodium hydroxide solution, in adding, while agitating, an alkali stabilizer, preservative and antibiotic solution, which is previously prepared in a sep. reactor, by mixing the aqueous solns. of benzalkonium chloride, disodium edetate and gentamicin sulfate, in subsequently adding a 10-20% sodium hydroxide solution, in adding xymedon at a

mass concentration of 1-10%, in adjusting, for the second time, the pH value to a required value by means of a sodium hydroxide solution and in sterilizing the thus produced del.

CC 63-6 (Pharmaceuticals)

REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L62 ANSWER 2 OF 19 ZCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 2

ACCESSION NUMBER: 2008:42992 ZCAPLUS Full-text

DOCUMENT NUMBER: 148:128281

TITLE: Nootropic medicinal agent

INVENTOR(S): Rasnetsov, Lev Davidovich; Shvartsman, Iakov

Yudelevich; Yashnova, Olga Konstantinovna; Melnikova,

Nina Borisovna; Petryakova, Olga Vladimirovna;

Gulyaev, Ivan Valeryevich

PATENT ASSIGNEE(S): Russia

SOURCE: PCT Int. Appl., 26pp.
CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: Russian

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PA'	PATENT NO.			KIN	KIND DATE			APPLICATION NO.					DATE				
	WO 2008004908			7.1	3.1 20000110			NO 2007 DU226					20070615				
WU	2000	0045	00		L/T		2000	0110		WO Z	007-	NU32	U		2007061		
	W:	ΑE,	AG,	AL,	AM,	ΑT,	AU,	ΑZ,	BA,	BB,	BG,	BH,	BR,	BW,	BY,	ΒZ,	CA,
		CH,	CN,	CO,	CR,	CU,	CZ,	DE,	DK,	DM,	DO,	DZ,	EC,	EE,	EG,	ES,	FI,
		GB,	GD,	GE,	GH,	GM,	GT,	HN,	HR,	HU,	ID,	IL,	IN,	IS,	JP,	KE,	KG,
		KM,	KN,	KP,	KR,	KZ,	LA,	LC,	LK,	LR,	LS,	LT,	LU,	LY,	MA,	MD,	ME,
		MG,	MK,	MN,	MW,	MX,	MY,	MZ,	NA,	NG,	NI,	NO,	NZ,	OM,	PG,	PH,	PL,
		PT,	RO,	RS,	SC,	SD,	SE,	SG,	SK,	SL,	SM,	SV,	SY,	TJ,	TM,	TN,	TR,
		TT,	TZ,	UA,	UG,	US,	UZ,	VC,	VN,	ZA,	ZM,	ZW					
	RW:	ΑT,	BE,	BG,	CH,	CY,	CZ,	DE,	DK,	EE,	ES,	FI,	FR,	GB,	GR,	HU,	IE,
		IS,	IT,	LT,	LU,	LV,	MC,	MT,	NL,	PL,	PT,	RO,	SE,	SI,	SK,	TR,	BF,
		ВJ,	CF,	CG,	CI,	CM,	GA,	GN,	GQ,	GW,	ML,	MR,	ΝE,	SN,	TD,	TG,	BW,
		GH,	GM,	KE,	LS,	MW,	MZ,	NA,	SD,	SL,	SZ,	TZ,	UG,	ZM,	ZW,	AM,	AZ,
		BY,	KG,	KZ,	MD,	RU,	TJ,	TM									
RU	2322	240			C1		2008	0420		RU 2	006-	1241	17		2	0060	705

PRIORITY APPLN. INFO.:

The invention relates to medicine, in particular to neurol, and psychiatry and can be used in the form of an agent for normalizing the physiol. and functional activity of the central nervous system of brain intellectual function. The inventive medicinal agent exhibits a nootropic activity and comprises dimephosphone in the form of an active substance. Said medicinal agent is embodied in the form of an aqueous solution and also comprises citric acid and lithium carbonate at the following component ratio: 5.0-30.0 mass% dimephosphone, 0.5-5.0 mass% lithium carbonate, 3.0-4.0 mass% citric acid, the rest up to 100% being deionized water. The medicinal agent in the form of a syrup has the following component ratio: 3.0-5.0 mass% dimephosphone, 1.0-1.2 mass% lithium carbonate, 4.0-5.0 mass% citric acid and 89-92.0 mass% sixtyfour percentage sugar syrup. A syrup having a high concentration of dimephosphone and a glycerin-containing syrup are also disclosed as the variants of the invention. Said medicinal agents have more physiol. pH values exhibited within an extended range of concns., are characterized by the high dilution stability of the solns. and have a delectable flavor, thereby easing the use thereof in a child treatment form.

RU 2006-124117 A 20060705

CC 63-6 (Pharmaceuticals)

Section cross-reference(s): 1

REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS

RECORD, ALL CITATIONS AVAILABLE IN THE RE FORMAT

L62 ANSWER 3 OF 19 ZCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 3

ACCESSION NUMBER: 2008:10448 ZCAPLUS Full-text

DOCUMENT NUMBER: 148:106213

TITLE: Pharmaceutical composition for treating burns and a

method for its production
INVENTOR(S): Rasnetsov, Lev Davidovich; Shvartsman, Iakov

Yudelevich: Yashnova, Olga Konstantinovna; Melnikova,

Nina Borisovna; Sorokin, Pavel Vladimirovich;

Zimnyakova, Olga Evgenyevna

PATENT ASSIGNEE(S): Russia

SOURCE: PCT Int. Appl., 25pp.

CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: Russian

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.			KIN	KIND DATE			- 2	APPLICATION NO.					DATE				
WO 2008002196			A1		2008	0103	1	WO 2007-RU327					20070615				
	W:	ΑE,	AG,	AL,	AM,	AT,	AU,	AZ,	BA,	BB,	BG,	BH,	BR,	BW,	BY,	BZ,	CA,
		CH,	CN,	CO,	CR,	CU,	CZ,	DE,	DK,	DM,	DO,	DZ,	EC,	EE,	EG,	ES,	FI,
		GB,	GD,	GE,	GH,	GM,	GT,	HN,	HR,	HU,	ID,	IL,	IN,	IS,	JP,	KE,	KG,
		KM,	KN,	ΚP,	KR,	ΚZ,	LA,	LC,	LK,	LR,	LS,	LT,	LU,	LY,	MA,	MD,	ME,
		MG,	MK,	MN,	MW,	MX,	MY,	MZ,	NA,	NG,	NI,	NO,	NZ,	OM,	PG,	PH,	PL,
		PT,	RO,	RS,	SC,	SD,	SE,	SG,	SK,	SL,	SM,	SV,	SY,	TJ,	TM,	TN,	TR,
		TT,	TZ,	UA,	UG,	US,	UZ,	VC,	VN,	ZA,	ZM,	ZW					
	RW:	AT,	BE,	BG,	CH,	CY,	CZ,	DE,	DK,	EE,	ES,	FI,	FR,	GB,	GR,	HU,	IE,
		IS,	IT,	LT,	LU,	LV,	MC,	MT,	NL,	PL,	PT,	RO,	SE,	SI,	SK,	TR,	BF,
		ВJ,	CF,	CG,	CI,	CM,	GA,	GN,	GQ,	GW,	ML,	MR,	NE,	SN,	TD,	TG,	BW,
		GH,	GM,	KE,	LS,	MW,	MZ,	NA,	SD,	SL,	SZ,	TZ,	UG,	ZM,	ZW,	AM,	AZ,
		BY,	KG,	ΚZ,	MD,	RU,	ТJ,	TM									
RU 2	2317	311			C1		2008	0227	1	RU 2	006-	1217	94		2	0060	619

PRIORITY APPLN. INFO.: RU 2006-121794 A 20060619 AB The invention relates to medicine, in particular to soft medicinal agents for external application (ointment, gels, emulsions, liniments) and can be used for treating thermal, solar and chemical burns of human beings and animals. The inventive pharmaceutical composition for treating burns is embodied in the form of a gel and contains an active substance N-(B-oxyethyl)-4.6dimethyldihydropyrimidone-2 (xymedon), a gel former, a moisture retaining agent and distilled water. The gel former can be embodied in the form of sodium, CM-cellulose, sodium alginates or the mixture thereof and the moisture retaining agent is embodied in the form of glycerin. In the other embodiments, addnl. to xymedon, the composition can contain an active substance in the form of silver nitrate or silver nitrate and sodium sulphacyl or levomycetin and succinic acid. The test have proved the high efficiency of said composition, which meets all the medical and biol. requirements of modern medicinal agents used for treating burns and wounds. The pharmaceutical composition is embodied in the form of a gel and can be used as highly efficient regenerating, wound-healing and micro-circulation improving means for treating infected burn wound.

CC 63-6 (Pharmaceuticals)

REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L62 ANSWER 4 OF 19 ZCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 4

ACCESSION NUMBER: 2008:1133434 ZCAPLUS Full-text
DOCUMENT NUMBER: 149:386541

TITLE: Antiviral medicine

INVENTOR(S): Rasnetsov, L. D.; Shvartsman, Ya. Yu.; Lyalina, I. K.

PATENT ASSIGNEE(S): Russia SOURCE: Russ.,

SOURCE: Russ., 4pp.
CODEN: RUXXE7
DOCUMENT TYPE: Patent

LANGUAGE: Russian

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
RU 2333753 PRIORITY APPLN. INFO.:	C1	20080920	RU 2007-107866 RU 2007-107866	20070302
PRIORITI APPEN. INFO.:			NO 2007-107000	20070302

AB The invention concerns the chemical pharmaceutical industry, particularly a medicine for treatment of viral disease, including HIV infection (AIDS and HIV related diseases). The antiviral medicine is a solution containing fullewene-polyhydropolyaminocaproic acid and DMSO in the following component content per ampoule: fullewene-polyhydropolyaminocaproic acid 50 mg, DMSO up to 0.5 mL. The medicine is intended for i.m. or i.v. (drip-feed) administration, and is dissolved in 20 mL of water for i.m. administration, while the variant for i.v. administration includes addnl. 20 mL of 0.9% sodium chloride solution per 1 ampoule. This provides an antiviral medicine for treatment of virus diseases including HIV infection.

- CC 63-6 (Pharmaceuticals)
- ST AIDS HIV virucide soln injection fullerene
 - T Fullerenes

Polvamides, biological studies

RL: PAC (Pharmacological activity); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)

(antiviral injection solution)

IT 25038-54-4D, Poly(6-aminocaproic acid), reaction product with

fullerenes, biological studies 99685-96-8, Fullerene 99685-96-8D, Fullerene, reaction products with poly(aminocaproic

acid) 131159-39-2, Fullerene

RL: PAC (Pharmacological activity); PEP (Physical, engineering or chemical process); TMU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)

(antiviral injection solution)

L62 ANSWER 5 OF 19 ZCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 5 ACCESSION NUMBER: 2008:1133379 ZCAPLUS Full-text

DOCUMENT NUMBER: 149:386539

TITLE: Antiviral medicine

INVENTOR(S): Rasnetsov, L. D.; Shvartsman, Ya. Yu.; Lyalina, I. K. PATENT ASSIGNEE(S): Russia

SOURCE: Russ., 4pp.
CODEN: RUXXE7

DOCUMENT TYPE: Patent LANGUAGE: Russian

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
RU 2333752	C1	20080920	RU 2007-107864	20070302
PRIORITY APPLN. INFO.:			RU 2007-107864	20070302

HIV-related diseases). The antiviral medicine is a 1% ointment containing fullerene-polyhydropolyaminocaproic acid as active substance and auxiliary substances of dimethylsulfoxide, water-free lanolin and Vaseline in the following amount (g): fullerene-polyhydropolyaminocaproic acid 1.0, dimethylsulfoxide 10.0, water-free lanolin 10.0, Vaseline up to 100.0. This provides an antiviral medicine for treatment of virus diseases, including HIV infection.

CC 63-6 (Pharmaceuticals)

ST HIV AIDS virucide ointment fullerene

IT Fullerenes

RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(antiviral ointment for AIDS)

IT 25038-54-4D, Poly(6-aminocaproic acid), reaction products with fullerenes, biological studies 99685-96-8, Fullerene

99685-96-8D, Fullerene, reaction products with poly(aminocaproic

acid) 131159-39-2, Fullerene RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL

(Biological study); USES (Uses)

(antiviral ointment for AIDS)

L62 ANSWER 6 OF 19 ZCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 6
ACCESSION NUMBER: 2008:1133377 ZCAPLUS Full-text

DOCUMENT NUMBER: 149:386538

TITLE: Antiviral medicine

INVENTOR(S): Rasnetsov, L. D.; Shvartsman, Ya. Yu.; Lyalina, I. K.

PATENT ASSIGNEE(S): Russia
SOURCE: Russ., 4pp.
CODEN: RUXXE7

DOCUMENT TYPE: Patent

LANGUAGE: Russian

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
RU 2333751	C1	20080920	RU 2007-107863	20070302
PRIORITY APPLN. INFO.:			RU 2007-107863	20070302

- AB The invention concerns the chemical pharmaceutical industry, particularly medicine for treatment of virus diseases, including HIV infection (AIDS and HIV-related diseases). The antiviral medicine is a suppository containing fullerene-polyhydropolyaminocaproic acid and auxiliary substances of dimethylsulfoxide, water-free lanolin or vegetable oil selected out of olive, peach, pumpkin seed oil, and a base selected out of M-35 or H-15 Witepsol, cacao butter, solid fat, in the following amount per one 2 g suppository: fullerene-polyhydropolyaminocaproic acid 5-20 mg, dimethylsulfoxide 50-200 mg, water-free lanolin or vegetable oil 20-100 mg, the rest being the base. This provides an antiviral medicine for treatment of virus diseases, including HIV infection.
- CC 63-6 (Pharmaceuticals)
- IT Fullerenes

RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (antiviral suppositories)

T 25038-54-4D, Poly(6-aminocaproic acid), reaction products with fullexenes, biological studies 99685-96-8D, Fullexene,

reaction products with poly(aminocaproic acid)
RL: MOA (Modifier or additive use); PAC (Pharmacological activity); THU
(Therapeutic use); BIOL (Biological study); USES (Uses)

(antiviral suppositories)

IT 99685-96-8, Fullerene 131159-39-2, Fullerene

RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(antiviral suppositories)

L62 ANSWER 7 OF 19 ZCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 7

ACCESSION NUMBER: 2008:171241 ZCAPLUS Full-text

DOCUMENT NUMBER: 148:222016

TITLE: Anti-viral agent for systemic application

INVENTOR(S): Rasnetsov, L. D.; Shvartsman, Ya. Yu.; Lyalina, I. K.

PATENT ASSIGNEE(S): Russia Russ., 5pp. SOURCE:

CODEN: RUXXE7

DOCUMENT TYPE: Patent

LANGUAGE: Russian

FAMILY ACC. NUM. COUNT: 1 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
RU 2316321	C1	20080210	RU 2006-121805	20060619
PRIORITY APPLN. INFO.:			RU 2006-121805	20060619
AB The invention pert.	ains to	the pharmac	centical industry, in	particular to

to agents for viral disease treatment including HIV infection (AIDS and HIV-associated diseases). The claimed agent is in the form of a solution for i.v. administering and contains per 1 ampoule 3 % concentrate of Fullevir (fullerenopolyaminocaproic acid sodium salt) 1 q and water for injection up to 1 mL. Addnl. it contains 0.9 % solution of sodium chloride or 10 % solution of human albumen in amount of 100 mL per 1 ampoule. This decreased viral load in lymphocytes and blood serum and increased amount of CD-4 cells.

63-6 (Pharmaceuticals)

L62 ANSWER 8 OF 19 ZCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 8

ACCESSION NUMBER: 2008:171240 ZCAPLUS Full-text

DOCUMENT NUMBER: 148:222015 TITLE: Anti-viral agent

INVENTOR(S): Rasnetsov, L. D.; Shvartsman, Ya. Yu.; Lyalina, I. K. PATENT ASSIGNEE(S):

Russia Russ., 6pp.

CODEN: RUXXE7 DOCUMENT TYPE: Patent

LANGUAGE: Russian

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

SOURCE:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
RU 2316320	C1	20080210	RU 2006-121806	20060619
PRIORITY APPLN. INFO.:			RU 2006-121806	20060619

AB The invention pertains to the pharmaceutical industry, in particular agents for viral disease treatment including HIV infection (AIDS and HIV-associated diseases). The claimed agent, in form of a suppository, contains per one 2 g suppository Fullevir (fullerenopolyaminocaproic acid sodium salt) 20 mg as active ingredient and ancillary substances such as propylene glycol 200 mg and balance: Vitepsol. This is an effective agent having no adverse influence on peripheral blood and body systems.

CC 63-6 (Pharmaceuticals)

L62 ANSWER 9 OF 19 ZCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 9 ACCESSION NUMBER: 2007:1061803 ZCAPLUS Full-text

DOCUMENT NUMBER: 147:330519

TITLE: A vaginal antimicrobial suppository

INVENTOR(S): Rasnetsov, Lev Davidovich; Shvartsman, Takov Yudelevitch; Lyalina, Irina Konstantinovna

PATENT ASSIGNEE(S): Zakrytoe Aktsionernoe Obschestvo "Intelpharm", Russia

SOURCE: PCT Int. Appl., 17pp.

CODEN: PIXXD2 DOCUMENT TYPE: Patent

LANGUAGE: Russian

FAMILY ACC. NUM. COUNT: 1 PATENT INFORMATION:

PATENT NO. KIND DATE APPLICATION NO. DATE WO 2007105984 A1 20070920 WO 2007-RU25 20070124 W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GO, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM RU 2318523 C2 20080310 RU 2006-107864 20060313 EE 2008-57 20070124 RU 2006-107864 A 20060313 WO 2007-RU25 W 20070124 EE 200800057 A 20081015 PRIORITY APPLN. INFO.:

- The invention relates to the chemical and pharmaceutical industry, in AB particular to producing a suppository antimicrobial agent which comprises iodine and can be used in clin. practice for treating inflammatory diseases of a female genital sphere. The inventive suppository antimicrobial agent contains iodine in the form of an active substance and a filler and is characterised in that the active substance is embodied in the form of an iodine-dimethyl-sulfoxide (DMSO) combination at a ratio of 1:(1-10) and comprises a liposol. base in the form of the filler, wherein the iodine content in the suppository ranges from 10 to 200 mg. The suppositories exhibit an extended antimicrobial spectrum.
- CC 63-6 (Pharmaceuticals)
- Section cross-reference(s): 1

REFERENCE COUNT: THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS 4 RECORD, ALL CITATIONS AVAILABLE IN THE RE FORMAT

L62 ANSWER 10 OF 19 ZCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 10 ACCESSION NUMBER: 2005:991188 ZCAPLUS Full-text

DOCUMENT NUMBER: 143:269083

TITLE: Method of production of C60 and C70 fullerenes and reactor for production of fullerene black

INVENTOR(S): Rasnetsov, L. D.; Shvartsman, Ya. Yu.; Lvalina, I. K.; Karnatsevich, V. L.; Kirillov, A. I.; Kaverin, B.

S.: Lopatin, M. A.

PATENT ASSIGNEE(S): Zakrytoe Aktsionernoe Obshchestvo "Fulleren-Tsentr", Russia; Institut Metalloorganicheskoi Khimii im. G. A.

Razuvaeva RAN

Russ., No pp. given SOURCE:

CODEN: RUXXE7

DOCUMENT TYPE: Pat.ent.

AB

LANGUAGE: Russian FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
RU 2259942	C2	20050910	RU 2003-127108	20030909
PRIORITY APPLN. INFO.:			RU 2003-127108	20030909

In pharmacol, fields, the method includes evacuation of a hermetic chamber, then filling it with helium. A voltage is supplied to a cathode and anode located resp. in a cathode lead-in and anode lead-in. The cathode lead-in moves in a longitudinal direction and the anode lead-in is immovable. After anode burning, its replacement is performed automatically from a rod loader. The anode and cathode are enclosed in metal casing with open ends, mounted coaxially relative to the electrodes. The casing is turnable and its longitudinal axis coincides with the axis of the upper flange and lower flange. The upper flange is provided with axle with piston for forcing fullerene-containing black to a storage chamber mounted on the lower flange. The black thus obtained is subjected to treatment in a Soxlet apparatus with aromatic solvent-toluene. To this end, use is made of excessive amts. of black relative to saturated solution of mixture of fullerenes in toluene. extract containing ≤95% C60 settles on the hot bottom of apparatus The solution above sediment is enriched with C70 ≤70%. Fullerence C60 and C70 are separated independently and in parallel in chromatog. columns using activated charcoal as immovable phase. Toluene or chlorobenzene is used as movable phase. The target product is crystallized and is addnl. cleaned by recrystallization or sublimation in vacuum, thus obtaining C60 at purity 99.9% and C70 at purity 99.5%. The result is enhanced reliability and facilitated method.

IC ICM C01B031-02

ICS B01D011-02; B01D015-08

CC 49-1 (Industrial Inorganic Chemicals)

production of fullerene black)

Section cross-reference(s): 76

ST fullerene black prodn reactor

IT Charcoal

RL: NUU (Other use, unclassified); PEP (Physical, engineering or chemical process); PIP (Physical process); PROC (Process); USES (USES) (activated; production of C60 and C70 fullerenes and reactor for

IT Liquid chromatography

Reactors

Sacrificial anodes

Solvent extraction

(production of C60 and C70 fullerenes and reactor for production of fullerene black)

IT Carbon black, preparation

RL: PUR (Purification or recovery); SPN (Synthetic preparation); PREP (Preparation)

(production of C60 and C70 fullerenes and reactor for production of fullerene black)

IT 7440-59-7, Helium, processes

RL: CPS (Chemical process); NUU (Other use, unclassified); PEP (Physical, engineering or chemical process); PROC (Process); USES (Uses)

(production of C60 and C70 fullerenes and reactor for production of fullerene black)

IT 108-88-3, Toluene, uses 108-90-7, Chlorobenzene, uses
RL: NUU (Other use, unclassified); USES (Uses)

(production of C60 and C70 fullerenes and reactor for production of fullerene black)

IT 7440-44-0P, Carbon, preparation 99685-96-8P, C60 Fullerene

115383-22-7P, C70 Fullerene

RL: PUR (Purification or recovery); SPN (Synthetic preparation); PREP

(Preparation)

(production of C60 and C70 fullerenes and reactor for production of fullerene black)

L62 ANSWER 11 OF 19 ZCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 11

ACCESSION NUMBER: 2005:498437 ZCAPLUS Full-text

DOCUMENT NUMBER: 143:45328

TITLE: Polyhexamethylenequanidine-containing noncorrosive

disinfecting detergent compositions

INVENTOR(S): Rasnetsov, L. D.; Gaiduchenva, G. M.; Shvartsman,

Ya. Yu.; Kozhevnikov, V. G.; Filonov, V. P.;

Gaiduchenya, A. V.; Rasnetsova, B. E.

PATENT ASSIGNEE(S): Zakrytoe Aktsionernoe Obshchestvo "Desko", Russia SOURCE: Russ., No pp. given

CODEN: BUXXE7

DOCUMENT TYPE: Patent.

LANGUAGE: Russian

FAMILY ACC. NUM. COUNT: 1 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
RU 2253669	C1	20050610	RU 2003-136055	2003121
PRIORITY APPLN. INFO.:			RU 2003-136055	2003121
AB A disinfecting det	ergent	composition	comprises a mixture of	a nonioni

surfactant (6.8-11.7), anionic surfactant (3.1-5.8), and cationic surfactant (0.5-1.0%), an active cleaning component (3.0-9.8), a polyhexamethylenequanidine derivative as a disinfectant (0.5-6.0), and a solvent (to 100%), the active cleaning component being a mixture of sodium CMcellulose, and sodium salts of phosphoric acid, sulfuric acid, and silicic acid, and the cationic surfactant being a quaternary ammonium compound Preferably, the nonionic surfactant is a mixture of Neonol, Syntanol, and wetting agent DB, the cationic surfactant is alkyldimethylbenzylammonium chloride or didecyldimethylammonium chloride, and the anionic surfactant is a mixture of Sulfanol and fatty alc. sulfates. The detergent compns. have improved anticorrosive properties, increased antibacterial and fungicidal activity, and can be used for cleaning of various surfaces (e.g. metal, glass) in medicine, food industry, engineering, and household.

ICM C11D001-86

ICS C11D001-62; C11D003-04; C11D003-48

46-6 (Surface Active Agents and Detergents) Section cross-reference(s): 63

L62 ANSWER 12 OF 19 ZCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 12

ACCESSION NUMBER: 2003:826903 ZCAPLUS Full-text

DOCUMENT NUMBER: 140:113683

TITLE: Method for preparing water-soluble amino acid

derivatives of fullerene

INVENTOR(S): Rasnetsov, L. D.; Shvartsman, Ya. Yu.; Lvalina, I.

> K.; Rasnetsova, B. E.; Karnatsevich, V. L.; Suvorova, O. N.; Kutyreva, V. V.; Shchupak, E. A.;

Bazvakina, N. L.: Makarov, S. G.

PATENT ASSIGNEE(S): Zakrytoe Aktsionernoe Obshchestvo "Desko", Russia

SOURCE: Russ., No pp. given

CODEN: RUXXE7 DOCUMENT TYPE: Pat.ent.

LANGUAGE: Russian

FAMILY ACC. NUM. COUNT: 1

5

PATENT INFORMATION:

```
PATENT NO.
                       KIND DATE APPLICATION NO. DATE
                       ----
    RU 2213049
                       C1 20030927 RU 2002-118286 20020708
RU 2002-118286 20020708
PRIORITY APPLN. INFO.:
    The organic chemical, chemical technol. The invention relates to the improved
     method for preparing water-soluble amino acids derivs. of fullerene that can
     be used in pharmacol, and microbiol. Invention describes method for preparing
     water-soluble amino acid derivs. of fullerene of the general formula (I):
     HC60NH(CH2)nC00-Kt+ wherein C60 is a fullerene ring; Kt+ is hydrogen atom,
     ammonium or alkaline metal cation; n = 1, 3, 5. Method involves interaction
     of fullerene with amino acid salt at heating and the following isolation of
     the end product. Compound of the general formula (II): is used as amino acid
     salt wherein R is CqH2q+1; m = 3, 4; q = 2-5; Y- is chemical element taken
     among (Va) or (VIa) groups of Mendelevev's periodic system. Then compound of
     the general formula (III): is prepared wherein R, Y, n, m have values given
     above that is subjected for the following reactions: in the case for preparing
     the end product of the general formula (I) wherein Kt+ is hydrogen atom method
     involves effect with acid solution and if Kt+ is ammonium or alkaline metal
     cation method involves effect with corresponding salt. Proposed method does
     not require the special equipment and can be carried out using the
     conventional chemical equipment that results to the simplified technol.
     process and reduced cost of the end product.
     ICM C01B031-02
ΙĊ
     ICS C07C229-06; C07F009-10; C07F009-66; C07F009-90; C07F009-94;
         C07F011-00
CC
    49-8 (Industrial Inorganic Chemicals)
ST
    water soluble amino acid deriv fullerene prepn
ΤТ
    Amino acids, reactions
     RL: RCT (Reactant); RACT (Reactant or reagent)
       (method for preparing water-soluble amino acid derivs. of fullerene
     Amino acids, reactions
     RL: RCT (Reactant); RACT (Reactant or reagent)
       (salts; method for preparing water-soluble amino acid derivs. of
        fullerene)
ΙT
    99685-96-8DP, Fullerene, amino acid derivs.
     RL: IMF (Industrial manufacture); PREP (Preparation)
        (method for preparing water-soluble amino acid derivs. of fullerene
L62 ANSWER 13 OF 19 ZCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 13
ACCESSION NUMBER:
                       2003:826902 ZCAPLUS Full-text
DOCUMENT NUMBER:
                       140:96298
TITLE:
                       Method for preparing water-soluble salts of amino acid
                        derivatives of fullerene
INVENTOR(S):
                       Rasnetsov, L. D.; Shvartsman, Ya. Yu.; Lyalina, I.
                       K.; Rasnetsova, B. E.; Karnatsevich, V. L.;
                        Suvorova, O. N.; Kutvreva, V. V.; Shchupak, E. A.;
                        Bazyakina, N. L.; Makarov, S. G.
PATENT ASSIGNEE(S):
                       Zakrytoe Aktsionernoe Obshchestvo "Desko", Russia
SOURCE:
                       Russ., No pp. given
                        CODEN: RUXXE7
DOCUMENT TYPE:
                       Patent
LANGUAGE:
                        Russian
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:
```

APPLICATION NO.

PATENT NO. KIND DATE

DATE

LANGUAGE:

Russian

RU 2213048 C1 20030927 RU 2002-118282 20020708 PRIORITY APPLN. INFO.: RU 2002-118282 The invention relates to the improved method for preparing water- soluble salts of amino acid derivs. of fullerene that can be used in medicine, pharmacol. and microbiol. Invention describes method for preparing watersoluble salts of amino acid derivs, of fullerene of the general formula HC60NH(CH2)nC00M wherein C60 is a fullerene ring; M is alkaline metal; n = 1, 3. 5. The method involves interaction of fullerene with amino acid salt in an organic solvent medium at heating and the following isolation of the end product. Interaction reaction is carried out in the presence of low-mol. polyalkylene oxide with mol. mass 150-400 Da. The invention provides reduced process time, and reduced manufacturing cost due to use of inexpensive raw materials ICM C01B031-02 ICS C07C229-06 CC 49-5 (Industrial Inorganic Chemicals) ST alkali metal amino acid salt fullerene deriv manuf ΙT IR spectra (method for preparing water-soluble salts of amino acid derivs. of fullerene) Polyoxyalkylenes, processes RL: CPS (Chemical process); PEP (Physical, engineering or chemical process); PROC (Process) (method for preparing water-soluble salts of amino acid derivs, of fullerene) Amino acids, preparation RL: IMF (Industrial manufacture); PREP (Preparation) (salts, alkali metal fullerese derivs.; method for preparing water-soluble salts of amino acid derivs. of fullerene) 99685-96-8P. Fullerene RL: IMF (Industrial manufacture); PREP (Preparation) (alkali metal amino acid salt derivs.; method for preparing water-soluble salts of amino acid derivs. of fullerene) 99685-96-8DP, Fullerene, alkali metal amino acid salt derivs. 645420-16-2P 645420-18-4P 645420-20-8P 645420-22-0P 645420-23-1P 645420-24-2P RL: IMF (Industrial manufacture); PREP (Preparation) (method for preparing water-soluble salts of amino acid derivs. of fullerene) 108-88-3, Toluene, uses RL: NUU (Other use, unclassified); USES (Uses) (method for preparing water-soluble salts of amino acid derivs. of fullerene) 6610-05-5, Sodium y-aminobutyrate 48047-10-5, Potassium s-aminocaproate RL: RCT (Reactant); RACT (Reactant or reagent) (method for preparing water-soluble salts of amino acid derivs. of fullerene) L62 ANSWER 14 OF 19 ZCAPLUS COPYRIGHT 2009 ACS on STN ACCESSION NUMBER: 2008:1519339 ZCAPLUS Full-text TITLE: Device for obtaining fullerene containing soot INVENTOR(S): Rasnetsov, L. D.; Shvartsman, Ya. Yu.; Karnatsevich, V. L.; Kirillov, A. I.; Kaverin, B. S. PATENT ASSIGNEE (S): ZAO "Fulleren-Tsentr", Russia Russ., 7pp. SOURCE: CODEN: RUXXE7 DOCUMENT TYPE: Patent

FAMILY ACC. NUM. COUNT: 1 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
RU 2341452	C1	20081220	RU 2007-113546	20070411
PRIORITY APPLN. INFO.:			RU 2007-113546	20070411

AB FIELD: chemical; electricity.SUBSTANCE: proposed device contains a cold airtight chamber filled with helium. On opposite walls of the camera casing 1 with the help of ports 4 and 5 anode current leads 6 and 7 are installed, in which electrodes 8 and 9 in the form of rods are placed in line with each other. Current leads 6 and 7 are connected to different power sources. Between rods 8 and 9 is placed a graphite electrode in the form of disk 10 with the formation of a discharge gap between them. Disk 10 is installed in a fixed position on the cathode current 11, which is placed on the upper flange of camera 2 and is connected to the elec. motor 12 for ensuring the possibility of the rotation of disk 10 on a plane parallel to the plane of flange 2. Burnt rods 8 and 9 are installed with the capability of moving in the discharge gap zone. On the outside of ports 4 and 5 are connected vacuum loaders of rods 13 each of which consists of cover 14, connected to its own port 4 and 5, vibration-layer 15, which contains the reserve rods, supply device, made, for example in the form of a closed chain for supply 16 toothpushers 17 with the capability of catching and moving rods 8 and 9, and intermittent drive 18. Cathode current lead 11 is supplied with knives 19 to prevent the possibility of outgrowths forming on disk 10. The lower flange 3 is connected to the soot accumulator 20.EFFECT: doubling the productivity of the device with continuous submission of the burnt rods to the zone of the discharge gap due to the organization of two arc processes in one chamber.l

49 (Industrial Inorganic Chemicals)

L62 ANSWER 15 OF 19 ZCAPLUS COPYRIGHT 2009 ACS on STN 2008:690830 ZCAPLUS Full-text ACCESSION NUMBER:

TITLE: Synthesis and properties of water-soluble fullerene

derivatives

AUTHOR(S): Suvorova, O. N.; Kutureva, V. V.; Baziakina, N. L.;

Karnatsevich, V. L.; Schupak, E. A.; Rasnetsov, L.

D.; Makarov, S. G.

CORPORATE SOURCE: Razuvaev Institute of Organometallic Chemistry of

Russian Academy of Sciences, Nizhnii Novgorod, 603900,

Russia

SOURCE: Hydrogen Materials Science and Chemistry of Carbon

Nanomaterials, International Conference, 9th,

Sevastopol, Ukraine, Sept. 5-11, 2005 (2005), 498-501. Editor(s): Schur, D. V.; Zaginaichenko, S. Yu.;

Veziroglu, T. Nejat. Association for Hydrogen Energy

in Ukraine: Kiev, Ukraine.

CODEN: 69KTNL

DOCUMENT TYPE: Conference LANGUAGE:

English/Russian

The processes of fullerene amino acids preparation by the direct addition of amino acid derivs. to $C60\Psi$ were investigated. Some technol. aspects of this reaction were studied using different phase-transfer catalysts, and new methods of fullerene amino acids production with quant. yields and their purification were suggested. Water-soluble derivs, were obtained using the method of 1,3-dipolar cycloaddn. of azomethine ylides via the decarboxilation of immonium salt derived from the condensation of sarcosine with Boc-protected amino ketone. The results show that both methods of fullerene functionalization can be successfully used for preparation of water-soluble fullerene derivs.

52 (Electrochemical, Radiational, and Thermal Energy Technology)

water soluble fullerene deriv optical property

IT INDEXING IN PROGRESS

IT IR spectroscopy

(synthesis and properties of water-soluble fullerene derivs.)

7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L62 ANSWER 16 OF 19 ZCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2008:690800 ZCAPLUS Full-text

X-ray rapid analysis of fullerene content in arc soot TITLE: AUTHOR(S): Kirillov, A. I.; Karnatsevich, V. L.; Rasnetsov, L. D. CORPORATE SOURCE: Institute of Organo-metallic Chemistry of RAS, Nizhny

Novgorod, 603095, Russia

Hydrogen Materials Science and Chemistry of Carbon SOURCE: Nanomaterials, International Conference, 9th,

Sevastopol, Ukraine, Sept. 5-11, 2005 (2005), 418-421. Editor(s): Schur, D. V.; Zaginaichenko, S. Yu.; Veziroglu, T. Nejat. Association for Hydrogen Energy

in Ukraine: Kiev, Ukraine.

CODEN: 69KTNL

DOCUMENT TYPE: Conference LANGUAGE: English/Russian

A method for the quant. X-ray anal. of fullerene-containing soot is developed. In the method, the samples are taken from different parts of the reactor with various content of fullereng. The concentration is determined by weighing the residue from Sokslet container. The measurement is carried out on (computerized) DRON-3M diffractometer with Cu-Kα radiation in step mode at narrow angle intervals under the peaks of fullerene and carbon. The merits of the method include: (1) no need to weigh a sample since only the total intensity of diffracted beam depends upon quantity of substance in the sample (d. of cell packing) as the ratio of intensities, being proportional to phase concns., remains constant, and (2) the time of anal. with subsequent processing of the result is about one hour which is by an order of magnitude less than in the above-mentioned techniques.

CC 52 (Electrochemical, Radiational, and Thermal Energy Technology)

ST arc soot fullerene X ray analysis

ΙT INDEXING IN PROGRESS

IT X-ray spectroscopy

(X-ray rapid anal. of fullerene content in arc soot)

L62 ANSWER 17 OF 19 ZCAPLUS COPYRIGHT 2009 ACS on STN ACCESSION NUMBER: 1996:746375 ZCAPLUS Full-text

DOCUMENT NUMBER: 126:24049

ORIGINAL REFERENCE NO.: 126:4831a,4834a

TITLE: Sorbent for radionuclide extraction INVENTOR(S):

Rasnetsov, Lev D.; Dyachkovskij, Fridrikh S.;

Tuzova, Alla M.; Rasnetsova, Betti B.; Fadeev, Vadim

V.; Kanakova, Olga A.; Zubkov, Aleksandr M.

PATENT ASSIGNEE(S): Aktsionernoe Obshchestvo Zakrytogo Tipa Aktsionernoe

Predprivatie "ring" Ltd, Russia

Russ. From: Izobreteniya 1996, (16), 168-170. SOURCE:

CODEN: RUXXE7

DOCUMENT TYPE: Pat.ent.

LANGUAGE: Russian

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO. KIND DATE APPLICATION NO. DATE

RU 2061540 C1 19960610 RU 1992-5059384 19920821 PRIORITY APPLN. INFO.: SU 1992-5059384 A 19920821

AB Title only translated.
IC ICM B01J020-00

IC ICM B01J020-00 ICS G21F009-12

CC 71-8 (Nuclear Technology)

L62 ANSWER 18 OF 19 ZCAPLUS COPYRIGHT 2009 ACS on STN ACCESSION NUMBER: 1996:147954 ZCAPLUS Full-text

DOCUMENT NUMBER: 124:192809

ORIGINAL REFERENCE NO.: 124:35339a,35342a

TITLE: Method for determining strontium radionuclides

INVENTOR(S): Spivakov, Boris Ya.; Petrukhin, Oleg M.; Rasnetsov,
Lev D.; Malofeeva, Galina I.; Danilova, Tatyana V.;

Tuzova, Alla M.; Rasnetsova, Betti E.
Aktsionernoe Obshchestvo Zakrytogo Tipa Aktsionernoe

Predprivatie "Ring" Ltd., Russia

SOURCE: Russ. From: Izobreteniya 1995, (17), 235.

CODEN: RUXXE7

DOCUMENT TYPE: Patent LANGUAGE: Russian

FAMILY ACC. NUM. COUNT: 1 PATENT INFORMATION:

PATENT ASSIGNEE(S):

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
RU 2037894	C1	19950619	RU 1992-5056410	19920727
PRIORITY APPLN. INFO.:			SU 1992-5056410 A	19920727

AB Title only translated.

IC ICM G21G004-00 ICS G01N030-06

CC 79-6 (Inorganic Analytical Chemistry) Section cross-reference(s): 71

L62 ANSWER 19 OF 19 WPIX COPYRIGHT 2009 THOMSON REUTERS on STN

ACCESSION NUMBER: 2004-735111 [72] WPIX <u>Full-text</u>
DOC. NO. CPI: C2004-258526 [72]

TITLE: Agent for inhibition of reproduction of enveloped

viruses, method for its preparing, pharmaceutical composition and method for inhibition of viral infections

DERWENT CLASS: B05

INVENTOR: LYALINA I K ; RASNETSOV L D; RASNETSOVA B E; SHVARTSAM I Y; SHVARTSMAN I Y; SHVARTSMAN L Y

PATENT ASSIGNEE: (DESK-R) DESKO STOCK CO; (LYAL-I) LYALINA I K; (RASN-I)
RASNETSOV L D; (RASN-I) RASNETSOVA B E; (SHVA-I)

SHVARTSMAN I Y; (RASN-I) RASNETSOVA B E; (SHVA-I)

SHVARTSMAN I Y; (RASN-I) RASNETSOV L
COUNTRY COUNT: 107

COUNTRY COUNT: 10/

PATENT INFORMATION:

PA:	TENT NO	KINI	DATE	WEEK	LA	PG	MAIN IPC
RU	2236852	C1	20040927	(200472)*	RU	0[0]	
WO	2004112804	A1	20041229	(200504)	RU		
EΡ	1645279	A1	20060412	(200626)	EN		
US	20060122276	A1	20060608	(200639)	EN		
BR	2004011679	A	20060829	(200659)	PT		

ΑU	2004249090	A1	20041229	(200660)	EN		
KR	2006017887	A	20060227	(200660)	KO		
CN	1819834	A	20060816	(200682)	z_{H}		
JP	2007522082	W	20070809	(200754)	JA	32	
TN	2006DN00326	P1	20070817	(200780)	EN		

APPLICATION DETAILS:

PATENT NO KIND	APPLICATION DATE
RU 2236852 C1	RU 2003-118500 20030623
AU 2004249090 A1	AU 2004-249090 20040531
BR 2004011679 A	BR 2004-11679 20040531
CN 1819834 A	CN 2004-80017167 20040531
EP 1645279 A1	EP 2004-748919 20040531
WO 2004112804 A1	WO 2004-RU208 20040531
EP 1645279 A1	WO 2004-RU208 20040531
US 20060122276 A1	WO 2004-RU208 20040531
BR 2004011679 A	WO 2004-RU208 20040531
KR 2006017887 A	WO 2004-RU208 20040531
JP 2007522082 W	WO 2004-RU208 20040531
US 20060122276 A1	US 2005-559681 20051206
KR 2006017887 A	KR 2005-724813 20051223
JP 2007522082 W	JP 2006-517009 20040531
IN 2006DN00326 P1	WO 2004-RU208 20040531
IN 2006DN00326 P1	IN 2006-DN326 20060118

FILING DETAILS:

PA	TENT NO	KIND			PAI	ENT	NO	
EP	1645279	A1	Based	on	WO	200	1112804	Α
	2004011679	A	Based				1112804	A
KR	2006017887	A	Based	on	WO	200	1112804	A
AU	2004249090	A1	Based	on	WO	200	1112804	A
JP	2007522082	W	Based	on	WO	200	1112804	A

JP 2007522082	W Based on	WO 2004112804	A
PRIORITY APPLN. INFO: INT. PATENT CLASSIF.:	RU 2003-118500	20030623	
MAIN:	A61K033-00		
SECONDARY:	A61K031-197; A61	K031-785; A61K038-55;	A61P031-18
IPC ORIGINAL:	A61K0031-185 [I,	C]; A61K0031-185 [I,0]; A61K0031-185
	[I,C]; A61K0031-	195 [I,A]; A61K0031-1	.97 [I,A];
	A61K0031-197 [I,	A]; A61K0031-197 [I,	A]; A61K0031-198
	[I,A]; A61K0031-	21 [I,C]; A61K0031-22	25 [I,A]; A61K0031-66
	[I,A]; A61K0031	-66 [I,C]; A61K0031-	74 [I,C]; A61K0031-74
	[I,C]; A61K0031	-74 [I,C]; A61K0031-7	785 [I,A];
	A61K0031-785 [I,	A]; A61K0031-785 [I,	A]; A61K0033-00 [I,A]
	; A61K0033-00 [I	,A]; A61K0033-00 [I,0]; A61K0038-55 [I,A]
	; A61K0038-55 [I	,A]; A61K0038-55 [I,A	A]; A61K0038-55 [I,C]
	; A61P0001-00 [I	,C]; A61P0001-16 [I, A	A]; A61P0031-00 [I,C]
	; A61P0031-00 [I	,C]; A61P0031-00 [I,0	C]; A61P0031-12 [I,A]
	; A61P0031-18 [I	,A]; A61P0031-18 [I,A	A]; A61P0031-18 [I,A]
	; A61P0031-22 [I	,A]	
IPC RECLASSIF.:	A61K0031-185 [I,	C]; A61K0031-197 [I,	A]; A61K0031-74 [I,C]
	; A61K0031-785 [I,A]; A61P0031-00 [I,	C]; A61P0031-18
	[I,A]		
ECLA:	A61K0031-197; A6	1K0031-785	

USCLASS NCLM: 514/567.000 NCLS: 977/738.000

JAP. PATENT CLASSIF .:

MAIN/SEC.: A61K0031-198; A61P0001-16; A61P0031-12; A61P0031-18;

A61P0031-22

FTERM CLASSIF.: 4C201; 4C206; 4C206/Aa01; 4C206/Aa02; 4C206/Fa53; 4C206/Ka08; 4C206/Ma04; 4C206/Ma37; 4C206/Ma51;

4C206/MA55; 4C206/NA14; 4C206/ZA75; 4C206/ZB33;

4C206/ZC55

BASIC ABSTRACT:

RU 2236852 C1 UPAB: 20050707

NOVELTY - Invention relates to the development of agent for inhibition of reproduction of enveloped viruses. Invention proposes the group of inventions combined by the general inventive project involving a method for preparing compounds, development of pharmaceutical compositions and methods for treatment using their, agent based on fullerene polycarboxylic anions for inhibition of activity of enveloped viruses in treatment of diseases caused by these viruses. Choice of such quantitative ratios of components and conditions for carrying out the reaction provide preparing products of poly-addition. In carrying out synthesis amount of amino acid has to exceed amount of fullerene by more 50 times. Invention relates also to a method for inhibition of reproduction of enveloped viruses in treatment of diseases caused by HIV/AIDS, herpes infections, viral hepatitis C. Invention provides preparing product that has unlimited solubility in water, necessary bioavailability, high effectiveness of effect on infected cells and low toxicity. The content of basic substance in the end product is 87%, not less. Process shows technological effectiveness and can be used in pharmaceutical industry. USE - Virology, pharmaceutical industry, pharmacy.

ADVANTAGE - Improved preparing method, improved inhibiting method, valuable medicinal properties of agent.5 cl MANUAL CODE: CPI: B10-B02; B10-C02; B10-J02; B14-A02A3; B14-A02A3; B14-A02A3; B10-G02; B10-J02; B10

B14-A02B1; B14-G01B

IN LYALINA I K ; RASNETSOV L D; RASNETSOVA B E; SHVARTSAM I Y; SHVARTSMAN I Y; SHVARTSMAN L Y

NOV NOVELTY - Invention relates to the development of agent for inhibition of reproduction of enveloped viruses. Invention proposes the group of inventions combined by the general inventive project involving a method for preparing compounds, development of pharmaceutical compositions and methods for treatment using their, agent based on fullerene polycarboxylic anions for inhibition of activity of enveloped viruses in treatment of diseases caused by these viruses. Choice of such quantitative ratios of components and conditions for carrying out the reaction provide preparing products of poly-addition. In carrying out synthesis amount of amino acid has to exceed amount of fullerene by more 50 times. Invention relates also to a method for inhibition of reproduction of enveloped viruses in treatment of diseases caused by HIV/AIDS, herpes infections, viral hepatitis C. Invention provides preparing product that has unlimited solubility in water, necessary bioavailability, high effectiveness of effect on infected cells and low toxicity. The content of basic substance in the end product is 87%, not less. Process shows technological effectiveness and can be used in pharmaceutical industry.

=> file registry
FILE 'REGISTRY' ENTERED AT 16:13:36 ON 02 JAN 2009
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERNS" FOR DETAILS.
COPYRIGHT (C) 2009 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 1 JAN 2009 HIGHEST RN 1092443-48-5 DICTIONARY FILE UPDATES: 1 JAN 2009 HIGHEST RN 1092443-48-5

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH July 5, 2008.

Please note that search-term pricing does apply when conducting ${\tt SmartSELECT}$ searches.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:


http://www.cas.org/support/stngen/stndoc/properties.html



chain nodes: 1 2 3 4 5 chain bonds: 1-2 2-3 3-4 3-5 exact/norm bonds: 3-4 3-5 exact bonds: 1-2 2-3

Connectivity: 2.2 E exact RC ring/chain
Match level: 1:CLASS 3:CLASS 4:CLASS 5:CLASS
Generic attributes: 2:
Saturation : Saturated

Uploading L19.str



ring nodes :

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20


ring bonds :

1-2 1-6 1-19 2-3 2-9 3-4 3-12 4-5 5-6 6-7 7-8 8-20 9-10 9-13 10-11 10-16 11-12 13-14 13-19 14-15 14-18 15-16 17-18 17-20 19-20 exact/norm bonds:

1-2 1-6 1-19 2-3 2-9 3-4 3-12 4-5 5-6 6-7 7-8 8-20 9-10 9-13 10-11 10-16 11-12 13-14 13-19 14-15 14-18 15-16 17-18 17-20 19-20

Match level :

1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom 11:Atom 12:Atom 13:Atom 14:Atom 15:Atom 16:Atom 17:Atom 18:Atom 19:Atom 20:Atom



1 2 3 4 5 7 8 9 10 11 12 chain bonds: 1-2 1-12 2-3 3-4 3-5 7-8 7-12 8-9 9-10 9-11 exact/norm bonds: 1-12 3-4 3-5 7-8 7-12 8-9 9-10 9-11 exact bonds:

1-2 2-3


chain nodes :

Connectivity :

21

```
2:2 E exact RC ring/chain
Match level:
1:CLASS 2:CLASS 3:CLASS 4:CLASS 5:CLASS 7:CLASS 8:CLASS 9:CLASS 10:CLASS
11:CLASS 12:Atom
Generic attributes:
2:
Saturation : Saturated
```

Element Count : Node 12: Limited C,C55



```
chain nodes:
1 2 3 4 5 7 8 9 10 11 12
chain bonds:
1-2 2-3 3-4 3-5 5-12 7-8 8-9 9-10 9-11 11-12
exact/norm bonds:
3-4 3-5 5-12 7-8 8-9 9-10 9-11 11-12
exact bonds:
1-2 2-3
```

Connectivity:
2:2 E exact RC ring/chain
Match level:
1:CLASS 2:CLASS 3:CLASS 4:CLASS 5:CLASS 7:CLASS 8:CLASS 9:CLASS 10:CLASS 11:CLASS 12:Atom
Generic attributes:
2:
Saturation : Saturated

Element Count :

Node 12: Limited C,C55

22

^{=&}gt; file zcaplus

FILE 'ZCAPLUS' ENTERED AT 16:13:40 ON 02 JAN 2009
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on SIN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS is strictly prohibited.

FILE COVERS 1907 - 2 Jan 2009 VOL 150 ISS 2 FILE LAST UPDATED: 1 Jan 2009 (20090101/ED)

ZCAplus now includes complete International Patent Classification (IPC) reclassification data for the third quarter of 2008.

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file contains CAS Registry Numbers for easy and accurate substance identification.
'OBI' IS DEFAULT SEARCH FIELD FOR 'ZCAPLUS' FILE

=> d stat que L26 L18 STR



Structure attributes must be viewed using STN Express query preparation. L19 $$\tt STR$$



Structure attributes must be viewed using STN Express query preparation. L21 $\,$ 260 SEA FILE=REGISTRY SSS FUL L18 AND L19 $\,$

L22	37	SEA	FILE=REGISTRY	SPE=ON	ABB=ON	PLU=ON	L21	AND	IDS/CI
L23	13	SEA	FILE=REGISTRY	SPE=ON	ABB=ON	PLU=ON	L22	AND	C60/CNS
L26	6	SEA	FILE=ZCAPLUS	SPE=ON	ABB=ON	PLU=ON	L23		


=> d stat que L43 L18 STR



Structure attributes must be viewed using STN Express query preparation. L19 $$\tt STR$$



Structure attributes must be viewed using STN Express query preparation. L21 260 SEA FILE-REGISTRY SSS FUL L18 AND L19 L39 STR



Structure attributes must be viewed using STN Express query preparation. ${\tt L40} \\ {\tt STR}$



```
Structure attributes must be viewed using STN Express guery preparation.
1.42
            15 SEA FILE=REGISTRY SUB=L21 SSS FUL L39 OR L40
L43
            15 SEA FILE=ZCAPLUS SPE=ON ABB=ON PLU=ON L42
=> s L26 or L43
           21 L26 OR L43
=> d stat que L9
            1 SEA FILE=REGISTRY SPE=ON ABB=ON PLU=ON 99685-96-8
          2246 SEA FILE=ZCAPLUS SPE=ON ABB=ON PLU=ON L2/D
T.3
           977 SEA FILE=ZCAPLUS SPE=ON ABB=ON PLU=ON L3 (L) PREP/RL
T. 4
L5
        207299 SEA FILE-ZCAPLUS SPE-ON ABB-ON PLU-ON AMINO ACID?/CW
L6
         15098 SEA FILE=ZCAPLUS SPE=ON ABB=ON PLU=ON L5 (L) RACT/RL
1.7
          4881 SEA FILE=ZCAPLUS SPE=ON ABB=ON PLU=ON L2 (L) RACT/RL
            30 SEA FILE=ZCAPLUS SPE=ON ABB=ON PLU=ON L6 AND L7
L8
             2 SEA FILE=ZCAPLUS SPE=ON ABB=ON PLU=ON L4 AND L8
L9
=> d stat que L12
             1 SEA FILE=REGISTRY SPE=ON ABB=ON PLU=ON 99685-96-8
L5
         207299 SEA FILE=ZCAPLUS SPE=ON ABB=ON PLU=ON AMINO ACID?/CW
L6
         15098 SEA FILE-ZCAPLUS SPE-ON ABB-ON PLU-ON L5 (L) RACT/RL
          4881 SEA FILE=ZCAPLUS SPE=ON ABB=ON PLU=ON L2 (L) RACT/RL
L7
            30 SEA FILE-ZCAPLUS SPE-ON ABB-ON PLU-ON L6 AND L7
L8
L11
        118009 SEA FILE=ZCAPLUS SPE=ON ABB=ON PLU=ON ?DIAMINO?/BI OR
               ?TRIAMINO?/BI OR DI AMINO?/BI OR ?TRI AMINO?/BI
             1 SEA FILE-ZCAPLUS SPE-ON ABB-ON PLU-ON L8 AND L11
=> s L26 or L43 or L9 or L12
           24 L26 OR L43 OR L9 OR L12
T.64
=> d ibib abs hitind hitstr L64 1-24
L64 ANSWER 1 OF 24 ZCAPLUS COPYRIGHT 2009 ACS on STN
ACCESSION NUMBER:
                       2008:1152155 ZCAPLUS Full-text
DOCUMENT NUMBER:
                        149:576255
TITLE:
                        Two Ih-symmetry-breaking C60 isomers stabilized by
                        chlorination
AUTHOR(S):
                        Tan, Yuan-Zhi; Liao, Zhao-Jiang; Oian, Zhuo-Zhen;
                        Chen, Rui-Ting; Wu, Xin; Liang, Hua; Han, Xiao; Zhu,
                        Feng; Zhou, Sheng-Jun; Zheng, Zhiping; Lu, Xin; Xie,
                       Su-Yuan; Huang, Rong-Bin; Zheng, Lan-Sun
CORPORATE SOURCE:
                       State Key Laboratory for Physical Chemistry of Solid
```

Surfaces and Department of Chemistry, College of

Chemistry and Chemical Engineering, Xiamen University,

Xiamen, 361005, Peop. Rep. China

SOURCE: Nature Materials (2008), 7(10), 790-794

CODEN: NMAACR; ISSN: 1476-1122

PUBLISHER: Nature Publishing Group

DOCUMENT TYPE: Journal LANGUAGE: English

AB One abiding surprise in fullerene science is that Ih-sym. buckminsterfullerene C60 (Ih-C60 or #1,812C60, the nomenclature specified by symmetry or by

Fowler's spiral algorithm) remains the sole C60 species exptl. available.

Setting it apart from the other 1,811 topol, isomers

(isobuckminsterfullerenes) is its exclusive conformity with the isolated-

pentagon rule, which states that stable fullerenes have isolated pentagons.

Although gas-phase existence of isobuckminsterfullerenes has long been suspected, synthetic efforts have yet to yield successful results. Here, the authors report the realization of two isobuckminsterfullerenes by chlorination

of the resp. C2v- and Cs-sym. C60 cages. These chlorinated species,

#1,809C60Cl8 and #1,804C60Cl12, were isolated in exptl. useful yields. Structural characterization by crystallog. unambiguously established the unique pentagon-pentagon ring fusions. These distinct structural features are

directly responsible for the regioselectivity observed in subsequent substitution of chlorines, and also render these unprecedented derivs. of C60

isomers important for resolving the long-standing puzzle of fullerene formation by the Stone-Wales transformation scheme.

CC 25-29 (Benzene, Its Derivatives, and Condensed Benzenoid Compounds) Section cross-reference(s): 22, 75, 78

IT 1082608-39-6P 1082608-40-9P 1082608-41-0P

RL: SPN (Synthetic preparation); PREP (Preparation) (preparation and regioselective substitution reactions of chlorinated fullerene-C60-C2v)

TT 1082608-41-0P


RL: SPN (Synthetic preparation); PREP (Preparation)

(preparation and regioselective substitution reactions of chlorinated fullerene-C60-C2v)

RN 1082608-41-0 ZCAPLUS

CN INDEX NAME NOT YET ASSIGNED

PAGE 1-A



 $\label{eq:page_2-A} \text{MeC-} \bigcup_{\text{CH}_2-\text{NH}}^{\text{CH}_2} \text{CH}_2 - \underbrace{\bigcup_{\text{NH}-\text{CH}_2-\text{OMe}}^{\text{OMe}}}_{\text{NH}-\text{CH}_2-\text{OMe}}$

REFERENCE COUNT: 30 THERE ARE 30 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L64 ANSWER 2 OF 24 ZCAPLUS COPYRIGHT 2009 ACS on STN ACCESSION NUMBER: 2007:418971 ZCAPLUS Full-text

DOCUMENT NUMBER: 147:188664

TITLE: Addition of bio-organic compounds on C60: A

semi-empirical investigation of its reactivity with

alvoine

AUTHOR(S): Ben Messaouda, Mhamed; Moussa, Fathi; Tangour, Bahoueddine; Szwarc, Henri; Abderrabba, Manef

CORPORATE SOURCE: Faculte de Pharmacie de Chatenay-Malabry, Universite

Paris XI, CNRS UMR 8612, Fr.
SOURCE: THEOCHEM (2007), 809(1-3), 153-159

CODEN: THEODJ; ISSN: 0166-1280

PUBLISHER: Elsevier B.V.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The thermodn. stability of C60(Glycine)n (where n = 1-4) has been studied by means of AM1 calcns. to determine the positions where glycine mols. are preferentially added onto [60]fullerene mol. This study is meant to get some insight into the results of syntheses of C60 derivs. with biol. activities.

CC 22-4 (Physical Organic Chemistry)
Section cross-reference(s): 34

TT 944383-07-7

RL: FMU (Formation, unclassified); PRP (Properties); RCT (Reactant); FORM (Formation, nonpreparative); RACT (Reactant or reagent)

(bis adduct, third glycine addition; semiempirical study of the energetics and regiochem. of addition reaction of C60 with glycine)

IT 944383-05-5

RL: FMU (Formation, unclassified); PRP (Properties); RCT (Reactant); FORM (Formation, nonpreparative); RACT (Reactant or reagent)

(lowest energy bis adduct, third glycine addition; semiempirical study of the energetics and regiochem. of addition reaction of C60 with glycine)

IT 944383-10-2

RL: FMU (Formation, unclassified); PRP (Properties); FORM (Formation, nonpreparative)

(lowest energy tetrakis adduct; semiempirical study of the energetics and regiochem. of addition reaction of C60 with glycine)

IT 944383-06-6

RL: FMU (Formation, unclassified); PRP (Properties); RCT (Reactant); FORM (Formation, nonpreparative); RACT (Reactant or reagent)

(lowest energy tris adduct, fourth glycine addition; semiempirical study of the energetics and regiochem. of addition reaction of C60 with glycine) 944393-09-9

RL: FMU (Formation, unclassified); PRP (Properties); FORM (Formation,

nonpreparative)

(tetrakis adduct; semiempirical study of the energetics and regiochem.

of addition reaction of C60 with glycine)

T 944383-08-8

RL: FMU (Formation, unclassified); PRP (Properties); RCT (Reactant); FORM (Formation, nonpreparative); RACT (Reactant or reagent)

(tris adduct, fourth glycine addition; semiempirical study of the energetics and regiochem. of addition reaction of C60 with glycine)


IT 944383-07-7

RL: FMU (Formation, unclassified); PRP (Properties); RCT (Reactant); FORM (Formation, nonpreparative); RACT (Reactant or reagent)

(bis adduct, third glycine addition; semiempirical study of the energetics and regiochem. of addition reaction of C60 with glycine)

RN 944383-07-7 ZCAPLUS

CN Glycine, N,N'-(9,59-dihydro[5,6]fullerene-C60-Ih-1,49-diyl)bis- (CA INDEX NAME)



IT 944383-05-5


RL: FMU (Formation, unclassified); PRP (Properties); RCT (Reactant); FORM (Formation, nonpreparative); RACT (Reactant or reagent)

(lowest energy bis adduct, third glycine addition; semiempirical study of the energetics and regiochem. of addition reaction of C60 with glycine)

RN 944383-05-5 ZCAPLUS

CN Glycine, N,N'-(9,32-dihydro[5,6]fullerene-C60-Ih-1,33-diyl)bis- (CA INDEX NAME)






- IT 944383-10-2
 - RL: FMU (Formation, unclassified); PRP (Properties); FORM (Formation, nonpreparative)
- (lowest energy tetrakis adduct; semiempirical study of the energetics and regiochem. of addition reaction of C60 with glycine)
- RN 944383-10-2 ZCAPLUS
 - CN Glycine, N,N',N'',N'''-(9,13,39,46-tetrahydro[5,6]fullerene-C60-Ih-1,14,38,58-tetrayl)tetrakis- (CA INDEX NAME)
- *** STRUCTURE DIAGRAM IS NOT AVAILABLE ***
- IT 944383-06-6
 - RL: FMU (Formation, unclassified); PRP (Properties); RCT (Reactant); FORM (Formation, nonpreparative); RACT (Reactant or reagent)
 - (lowest energy tris adduct, fourth glycine addition; semiempirical study

- of the energetics and regiochem. of addition reaction of C60 with glycine)
- RN 944383-06-6 ZCAPLUS
- CN Glycine, N,N',N''-(13,39-dihydro[5,6]fullerene-C60-Ih-1,14,38(9H)triyl)tris- (CA INDEX NAME)

PAGE 1-A





PAGE 3-A

HO2C-CH2-NH

IT 944383-09-9

RL: FMU (Formation, unclassified); PRP (Properties); FORM (Formation, nonpreparative)

(tetrakis adduct; semiempirical study of the energetics and regiochem. of addition reaction of C60 with glycine)

RN 944383-09-9 ZCAPLUS


CN Glycine, N,N',N'',N'''-(9,16,25,49-tetrahydro[5,6]fullerene-C60-Ih1,17,24,59-tetrayl)tetrakis- (CA INDEX NAME)

PAGE 1-A



PAGE 1-B

— CO2H



PAGE 2-B

-- co2H


IT 944383-08-8

RL: FMU (Formation, unclassified); PRP (Properties); RCT (Reactant); FORM (Formation, nonpreparative); RACT (Reactant or reagent) (tris adduct, fourth glycine addition; semiempirical study of the energetics and regiochem. of addition reaction of C60 with glycine)

RN 944383-08-8 ZCAPLUS

CN Glycine, N,N',N''-(32,42-dihydro[5,6]fullerene-C60-Ih-1,33,41(9H)-triyl)tris- (CA INDEX NAME)





NH-CH2-CO2H

PAGE 3-A

REFERENCE COUNT: 46 THERE ARE 46 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L64 ANSWER 3 OF 24 ZCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2007:365074 ZCAPLUS Full-text

DOCUMENT NUMBER: 147:10174

TITLE: Fullerene-derivatized amino acids: synthesis, characterization, antioxidant properties, and

solid-phase peptide synthesis

AUTHOR(S): Yang, Jianzhong; Alemany, Lawrence B.; Driver, Jonathan; Hartgerink, Jeffrey D.; Barron, Andrew R.

CORPORATE SOURCE: Richard E. Smallev Institute for Nanoscale Science and

Technology, The Institute of Biosciences and Bioengineering, and Center for Biological and Environmental Nanotechnology, Rice University,

Houston, TX, 77005, USA

SOURCE: Chemistry-A European Journal (2007), 13(9), 2530-2545

CODEN: CEUJED; ISSN: 0947-6539
PUBLISHER: Wiley-VCH Verlag GmbH & Co. KGaA

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 147:10174

A series of [60]fullerene-substituted phenylalanine (Baa) and lysine derivs. have been prepared by the condensation of 1,2-(4'-oxocyclohexano)fullerene with the appropriately protected (4-amino)phenylalanine and lysine, resp. Conversion of the imine to the corresponding amine was achieved by di-acid catalyzed hydroboration. The reduction of the imine was not accompanied by hydroboration of the fullerene cage. The [70]fullerene phenylalanine derivative has also been prepared as have the di-amino acid derivs. The compds. were characterized by MALDI-TOF mass spectrometry, UV/Vis spectroscopy, and cyclic voltammetry. 1H and 13C NMR spectroscopy allowed the observation of diastereomers. Fullerene-substituted peptides may be synthesized on relatively large scale by solid-phase peptide synthesis. The presence of the C60-substituted amino acid in a peptide has a significant effect on the secondary structures and self-assembly properties of peptides as compared to the native peptide. The antioxidant assay of Baa and a Baaderived anionic peptide was determined to be significantly more potent than Trolox.

CC 34-3 (Amino Acids, Peptides, and Proteins) Section cross-reference(s): 22

IT Amino acids, reactions

RL: RCT (Reactant); RACT (Reactant or reagent)

(N-[(fluorenylmethoxy)carbonyl]; preparation and antioxidant properties of fullerene-derivatized amino acids via condensation of fullerene ketone with amino acids followed by reduction, and their use in solid-phase synthesis of fullerene-peptide conjugates)

IT Amino acids, preparation

RL: PRP (Properties); RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(fullerene-derivatized; preparation and antioxidant properties of fullerene-derivatized amino acids via condensation of fullerene ketone with amino acids followed by reduction, and their use in solid-phase synthesis of fullerene-peptide conjugates)

99685-96-8, C60 Fullerene

RL: PRP (Properties); RCT (Reactant); RACT (Reactant or reagent)

(preparation and antioxidant properties of fullerene-derivatized amino

acids

via condensation of fullerene ketone with amino acids followed by reduction, and their use in solid-phase synthesis of fullerene-peptide conjugates)


99685-96-8, C60 Fullerene

RL: PRP (Properties); RCT (Reactant); RACT (Reactant or reagent) (preparation and antioxidant properties of fullerene-derivatized amino acids

via condensation of fullerene ketone with amino acids followed by reduction, and their use in solid-phase synthesis of fullerene-peptide conjugates)

RN 99685-96-8 ZCAPLUS

CN [5,6]Fullerene-C60-Ih (CA INDEX NAME)



REFERENCE COUNT: 33 THERE ARE 33 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L64 ANSWER 4 OF 24 ZCAPLUS COPYRIGHT 2009 ACS on STN ACCESSION NUMBER: 2006:960378 ZCAPLUS Full-text

DOCUMENT NUMBER: 146:371879

TITLE: Gene delivery by aminofullerenes: structural requirements for efficient transfection

AUTHOR(S): Isobe, Hiroyuki; Nakanishi, Waka; Tomita, Naoki; Jinno, Shigeki; Okayama, Hiroto; Nakamura, Eiichi

Department of Chemistry and ERATO (JST), The University of Tokyo, Hongo, Bunkyo-ku, Tokyo,

113-0033, Japan

SOURCE: Chemistry-An Asian Journal (2006), 1(1-2), 167-175

CODEN: CAAJBI; ISSN: 1861-4728

PUBLISHER: Wiley-VCH Verlag GmbH & Co. KGaA

DOCUMENT TYPE: Journal

CORPORATE SOURCE:

LANGUAGE: English

OTHER SOURCE(S): CASREACT 146:371879

A series of aminofullerenes that share a common structural motif have been synthesized and subjected to a systematic investigation of structure activity relationship regarding their ability for transient transfection and cytotoxicity. DNA-binding tests indicated that any water-soluble fullerenebearing amino group would bind to double-stranded DNA. For these mols, to be effective transfection reagents, however, they require addnl. structural features. First, the mol. must be capable of producing submicrometersized fullerene/DNA aggregates that can be internalized into mammalian cells through endocytosis. Second, the mol. must be capable of releasing DNA as the aggregates are transferred into the cytoplasm. This can be achieved in at least two ways: by loss of the DNA-binding amino groups from the fullerene core, and by transformation of the amino groups to neutral groups such as amides. The screening expts. led us to identify the best reagent, a tetrapiperidinofullerene, that can be synthesized in two steps from fullerene, piperazine, and mol. oxygen, and that is more efficient at transfection than a commonly used lipid-based transfection reagent.

CC 1-3 (Pharmacology)

T7 -44-3P, Spermine 110-60-1P, 1,4-Butanediamine 113-00-8P, Guanidine 124-20-9P, Spermine 144487-61-6P 169477-76-3P 169477-77-4P 18923-4B-0P 226420-73-1P 271785-61-6P 271785-65-0P 312773-17-4P 312773-18-5P 312773-19-6P 312773-20-9P 312773-22-1P 312773-23-2P 312773-24-3P 407617-27-0P 854752-05-9P 932025-64-4P 932025-65-5P 932025-66-6P 932025-67-7P 932025-68-8P 932025-64-4P 932025-65-5P 932025-71-3P RI: PAC (Pharmacological activity); PKT (Pharmacokinetics); SPN (Synthetic preparation); TBU (Therapeutic use); BIOL (Biological study); PREP

(gene delivery by aminofullerenes and structural requirements for efficient transfection)

IT 226420-73-1P 407617-27-0P

(Preparation); USES (Uses)

RL: PAC (Pharmacological activity); PKT (Pharmacokinetics); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)


(gene delivery by aminofullerenes and structural requirements for efficient transfection)

RN 226420-73-1 ZCAPLUS

CN Glycine, N-[3-(dimethylamino)propyl]-N-methyl-, 1,1'-(5',5''-hexano-3'H,3''H-dicyclopenta[1,9:13,14][5,6]fullerene-C60-Ih-3',3''-divl) ester (CA INDEX NAME)

PAGE 1-A

PAGE 2-A



RN 407617-27-0 ZCAPLUS


CN Glycine, N-[3-(dimethylamino)propyl]-N-methyl-,
[15,7'R,8'R,9S,9'R,135,13'R,145]-8',9'-dihydro-7',13'-dimethoxy-11',11'dimethyl-5',5''-(ethano! 4,5]-endo-[1,3]dioxoloethano)-3'H,3''Hdicyclopenta[1,9:13,14][5,6]fullerene-C60-Ih-3,3''-diyl ester (CA INDEX NAME)


Absolute stereochemistry.


PAGE 1-C















PAGE 4-A



REFERENCE COUNT: 71 THERE ARE 71 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L64 ANSWER 5 OF 24 ZCAPLUS COPYRIGHT 2009 ACS on STN ACCESSION NUMBER: 2006:162438 ZCAPLUS Full-text

DOCUMENT NUMBER: 145:455231

TITLE: Synthesis of fullerene-glycine derivative

AUTHOR(S): Jiang, Guichang; Zheng, Qixin

CORPORATE SOURCE: Department of Biology, Huazhong University of Science and Technology, Wuhan, 430074, Peop. Rep. China

SOURCE: Huagong Xinxing Cailiao (2005), 33(8), 24-26, 30 CODEN: HXCUA4; ISSN: 1006-3536

PUBLISHER: Huagong Xinxing Cailiao Bianjibu

DOCUMENT TYPE: Journal LANGUAGE: Chinese

OTHER SOURCE(S): CASREACT 145:455231

AB A novel fullerene-glycine derivative was synthesized by means of organic chemical It is soluble in polar solvents such as water, DMSO and THF et al. The product was characterized by FTIR, 1H-NMR and TEM. TEM anal. showed that it presents an ideal spherical shape in water with an average particle diameter of about 18mm. The in vitro antitumor activity of the novel derivative has been tested and the result showed that the novel derivative exhibited better antitumor activity in vitro against bone tumor cells. The in vitro antitumor activity of the novel derivative were related to the derivative concentration, and were also dependent on the power of the light irradiation The antitumor mechanism of the derivative was studied.

CC 34-2 (Amino Acids, Peptides, and Proteins)

Section cross-reference(s): 1

II 645420-16-2P, N-([5,6]Fulleren-C60-yl)glycine monosodium salt RL: PAC (Pharmacological activity); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)

(preparation of N-([5,6]fulleren-C60-yl)glycine sodium salt and study of

its activity as anticancer agent)

IT 645420-16-2P, N-([5,6]Fulleren-C60-yl)glycine monosodium salt RL: PAC (Pharmacological activity); SPN (Synthetic preparation); BIOL (Biological study); PREF (Preparation)

 $(preparation\ of\ N-([5,6]fulleren-C60-y1)glycine\ sodium\ salt\ and\ study\ of\ its$

activity as anticancer agent)

RN 645420-16-2 ZCAPLUS

CN Glycine, N-[5,6]fulleren-C60-Ih-1(?H)-yl-, monosodium salt (9CI) (CA INDEX NAME)

CM 1

CRN 645420-15-1 CMF C62 H63 N O2

*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

L64 ANSWER 6 OF 24 ZCAPLUS COPYRIGHT 2009 ACS on STN ACCESSION NUMBER: 2005:1298795 ZCAPLUS Full-text

DOCUMENT NUMBER: 144:260374

TITLE: Nonviral Gene Delivery by Tetraamino Fullerene AUTHOR(S): Isobe, Hirovuki; Nakanishi, Waka; Tomita, Naoki; Jinno, Shigeki; Okayama, Hiroto; Nakamura, Eiichi

CORPORATE SOURCE: Department of Chemistry and Department of Biochemistry and Molecular Biology (Graduate School of Medicine), University of Tokyo, Tokyo, 113-0033, Japan

SOURCE: Molecular Pharmaceutics (2006), 3(2), 124-134 CODEN: MPOHBP: ISSN: 1543-8384

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal LANGUAGE: English

CASREACT 144:260374 OTHER SOURCE(S):

A fullerene derivative bearing two diamino side chains binds to a plasmid vector DNA, either 4 or 40 kbp in size, delivers it to mammalian cells on incubation, and leads to expression of the encoded gene either transiently or stably. The initial physicochem, investigations upon DNA-binding and protective properties of various fullerene compds. against nuclease led us to identify the tetraamino fullerene as an ideal candidate to probe the new concept of fullerene-mediated gene delivery to mammalian cells. Studies on transient and stable transfection of COS-1 cells using green fluorescent protein and luciferase reporter genes revealed several useful properties of the fullerene transfection as compared with the conventional lipid-based transfection method, including much higher efficiency of stable transfection and ability to transfect confluent cells. Chemical and biol. studies suggested that the cell uptake of the fullerene/DNA complex takes place by an endocytosis mechanism and that the DNA internalized by endosomes is protected by the fullerene against enzymic digestion. The stiffness of the fullerene/DNA complex may play some role in the success of the fullerene method.

- 63-5 (Pharmaceuticals)
- 226420-73-1DP, complex with DNA

RL: BSU (Biological study, unclassified); PRP (Properties); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(nonviral gene delivery by tetraamino fullerene)

226420-73-1P

RL: PRP (Properties); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses) (nonviral gene delivery by tetraamino fullerene)

226420-73-1DP, complex with DNA


RL: BSU (Biological study, unclassified); PRP (Properties); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation): USES (Uses)

(nonviral gene delivery by tetraamino fullerene)


- 226420-73-1 ZCAPLUS RN
- CN Glycine, N-[3-(dimethylamino)propyl]-N-methyl-,

1,1'-(5',5''-hexano-3'H,3''H-dicyclopenta[1,9:13,14][5,6]fullerene-C60-Ih-3',3''-diyl) ester (CA INDEX NAME)

$$\begin{array}{c} \text{Me} \\ \text{CH}_2 - \text{NMe}_2 \end{array}$$



- IT 226420-73-1P
 - RL: PRP (Properties); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
- (nonviral gene delivery by tetraamino fullerene)
- RN 226420-73-1 ZCAPLUS
- CN Glycine, N-[3-(dimethylamino)propyl]-N-methyl-, 1,1'-(5',5''-hexano-3'H,3''H-dicyclopenta[1,9:13,14][5,6]fullerene-C60-Ih-3',3''-diyl) ester (CA INDEX NAME)



REFERENCE COUNT:

THERE ARE 69 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L64 ANSWER 7 OF 24 ZCAPLUS COPYRIGHT 2009 ACS on STN ACCESSION NUMBER: 2005:760228 ZCAPLUS Full-text

69

DOCUMENT NUMBER: 143:341234

Fractal Behavior of Functionalized Fullerene TITLE:

Aggregates. I. Aggregation of Two-Handed Tetraaminofullerene with DNA

AUTHOR(S): Ying, Oicong; Zhang, Jun; Liang, Dehai; Nakanishi, Waka; Isobe, Hiroyuki; Nakamura, Eiichi; Chu, Benjamin

CORPORATE SOURCE: Department of Chemistry, Stony Brook University, Stony Brook, NY, 11794-3400, USA

SOURCE: Langmuir (2005), 21(22), 9824-9831

AR

CODEN: LANGD5; ISSN: 0743-7463 PUBLISHER: American Chemical Society DOCUMENT TYPE: Journal

LANGUAGE: English In tris-buffered saline (TBS) with a trace of DMF, the homoaggregation process of a functionalized fullerene, the two-handed tetraaminofullerene (TH), and the heteroaggregation process (complex formation) of TH with DNA (pGL3-control plasmid) were studied dynamically by using a combination of static and dynamic laser light scattering measurements. Fractal behavior was investigated in the aggregation process of both TH homoaggregates and TH-DNA heteroaggregates. The stability of aggregates in solution depends on the molar concentration ratio RM, defined as the molar ratio of moles of TH to moles of the DNA base pair. Higher RM values resulted in lower aggregate stability. The transition of the fractal dimension (Df) in TH homoaggregation by rapidly mixing 3.78 µM TH with an equal volume of the blank buffer was found to vary from a value of 1.46 to 2.02. Dynamic light scattering results revealed that, in the aggregation process, the change in the size distribution of aggregates with time could be related to a Df transition. In the Df transition region, the size distribution of homoaggregates displayed a drastic change from a singlemode distribution to a bimodal distribution, which clearly suggested a restructuring process with the formation of large aggregates. When the aggregation process finally reached equilibrium, Df = 2.02, the size of the homoaggregates had a single mode but a broad distribution. However, TH-DNA heteroaggregation showed a Df transition from 1.58 to 1.7, but over a shorter time range of less than 5 min. Then, the Df value fluctuated in the range of 1.7 and finally reached an equilibrium value of Df \approx 1.78, which was independent of molar concentration There are two main action forces involved in the heteroaggregation process: van der Waals forces and attractive electrostatic forces, with the latter one being stronger and faster than that of the former. Therefore, a two-step action could occur in the heteroaggregation process. In the beginning of mixing, the attractive electrostatic forces dictated the aggregation process, and then van der Waals forces also got involved in the entire aggregation process. By using an initial concentration of 3.78 µM each and RM = 1, TH-DNA heteroaggregates showed more stable solution behavior than the homoaggregates. The lower Df value of the heteroaggregates could be related to a looser compact structure. Results from SEM also disclosed the different textures between TH homoaggregates and TH-DNA heteroaggregates; the former had a more dense

6-2 (General Biochemistry)

packing than the latter one.

IT 226420-73-1

> RL: BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)

(fractal behavior of functionalized fullerene aggregates and aggregation of two-handed tetraaminofullerene with DNA)

226420-73-1 IT

CN


RL: BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)

(fractal behavior of functionalized fullerene aggregates and aggregation of two-handed tetraaminofullerene with DNA)

226420-73-1 ZCAPLUS

Glycine, N-[3-(dimethylamino)propyl]-N-methyl-, 1,1'-(5',5''-hexano-3'H,3''H-dicyclopenta[1,9:13,14][5,6]fullerene-C60-Ih-3',3''-div1) ester (CA INDEX NAME)

$$\begin{array}{c} \text{Me} \\ \text{CH}_2 - \text{N}_{\text{L}} \text{ (CH}_2) \text{ } 3 - \text{NMe} \text{ } 2 \end{array}$$



REFERENCE COUNT:

THERE ARE 29 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L64 ANSWER 8 OF 24 ZCAPLUS COPYRIGHT 2009 ACS on STN ACCESSION NUMBER: 2005:554174 ZCAPLUS Full-text

29

DOCUMENT NUMBER: 144:331668

TITLE: Synthesis and solubility of 6-aminohexanoic acid and 2-aminoethanesulfonic acid C60 adducts

AUTHOR(S): Liu, Xu-Feng; Guan, Wen-Chao; Cheng, Zhen-Xian CORPORATE SOURCE: Department of Chemistry, Huazhong University of Science and Technology, Wuhan, 430074, Peop. Rep.

China

SOURCE: Youji Huaxue (2005), 25(6), 741-744 CODEN: YCHHDX: ISSN: 0253-2786

PUBLISHER: Youji Huaxue Bianjibu

DOCUMENT TYPE: Journal LANGUAGE: Chinese

OTHER SOURCE(S): CASREACT 144:331668

Description of water soluble fullerenes (C60) derivs. is meaningful to biol. Study of fullerenes. Amination reaction of amino-acid with C60 led to water soluble amino-acid C60 derivs. Reaction of C60 with excess of NH2(CH2)5COOM or NH2(CH2)2SO3Na (molar ratio is 1: 10) at 80 °C for 24 h afforded main amino-acid C60 adducts with addition degree of 5 and 4, resp. The yields based on the C60 added were 30% and 28%, resp. The addition degree was influenced by the length of hydrocarbon chain of amino-acid and precipitation of C60 adducts from the reactant. C60(NH(CH2)5CO0H)5H5 (I) and C60(NHCH2CH2SO3H)4H4 (II) were further purified by silica column chromatog. and characterized by IH NMR, 13C NMR, IR, FAB-MS spectra and elemental anal. The solubility of II was less pH dependent. The solubility of I in water at different pH was measured by the spectrophotometric method, exhibiting solubility of 71.81 mgmL-1 (pH = 10.25), 23.68 mgmL-1 (pH = 7) and 10.12 mgmL-1 (pH = 3.36). The s value of II at 272.8 nm was 3.43×104 Lnog-1-cm-1.

CC 34-2 (Amino Acids, Peptides, and Proteins)

IT 880763-63-3P 880763-66-6P

RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation) (synthesis and solubility of 6-aminohexanoic acid and 2-aminoethanesulfonic acid C60 adducts)

IT 880763-63-3P

RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation) (synthesis and solubility of 6-aminohexanoic acid and 2-aminoethanesulfonic acid C60 adducts)


RN 880763-63-3 ZCAPLUS

CN Hexanoic acid, 6,6',6'',6''',6'''',6''''-([5,6]fullerene-C60-Ih-pentaylpentaimino)pentakis- (9CI) (CA INDEX NAME)

CM

CRN 880763-62-2 CMF C90 H115 N5 O10

CCT TDS



5 D1-NH-(CH2)5-CO2H

DOCUMENT NUMBER: 141:388621

TITLE: Agent for inhibition of reproduction of enveloped viruses, method for its preparing, pharmaceutical

> composition and method for inhibition of viral infections

PATENT ASSIGNEE(S): Zakrytoe Aktsionernoe Obshchestvo "Desko", Russia SOURCE:

Russ., No pp. given

CODEN: RUXXE7

DOCUMENT TYPE: Pat.ent. LANGUAGE: Russian


FAMILY ACC. NUM. COUNT: 1 PATENT INFORMATION:

	PATENT NO.						DATE			APPLICATION NO.							
	2236852						20040927			RU 2003-118500							
AU	2004249090				A1 200412			1229	AU 2004-249090				20040531				
CA	2530004				A1 20041229			CA 2004-2530004					20040531				
WO					A1 20041229			WO 2004-RU208				20040531					
											, BG,						
											, EC,						
											, JP,						
											, MK,						
											, SD,						
											, vc,						,
	RW:										, SL,						AM.
											, BE,						
											LU.						
											, GA,						
			TD,		,	,	O. ,	,	,	0.,	,,	01.7	02,	,	,	,	,
EP	1645				A1		2006	0412		EP	2004-	7489	19		2	0040	531
											, IT,						
											, HU,			,	02,	110,	,
CN	1819				A						2004-				2	0040	531
	2004										2004-						
.TP	2007	5220:	R 2		т						2006-					0040	
	JP 2007522082 US 20060122276								US 2005-559681								
	2006															0060	
	IORITY APPLN. INFO.:				-		20070817			IN 2006-DN326 RU 2003-118500							
	101111111111111111111111111111111111111										2003-					0030	
munn o	UED COUDON (C)				143.01	D 7 FF	1 41 .	2000			2004-		~		. 2	0040	J J I

OTHER SOURCE(S): MARPAT 141:388621

Agent for inhibition of reproduction of enveloped viruses, method for its preparing, pharmaceutical composition and method for inhibition of viral infections are disclosed. The invention relates to the development of agent for inhibition of reproduction of enveloped viruses. The invention proposes the group of inventions combined by the general inventive project involving a method for preparing compds., development of pharmaceutical compns. and methods for treatment using those compds. and compns., agent based on fullerene polycarboxylic anions for inhibition of the activity of enveloped viruses in treatment of the diseases caused by these viruses. Choice of such quant. ratios of components and conditions for carrying out the reaction provide preparing products of poly-addition In carrying out synthesis amount of amino acid has to exceed amount of fullerene by more 50 times. The invention relates also to a method for inhibition of reproduction of enveloped viruses in treatment of diseases caused by HIV/AIDS, herpes infections, viral hepatitis C. Invention provides preparing product that has unlimited solubility in water, necessary bioavailability, high effectiveness on infected cells and low toxicity. The content of basic substance in the end product is 87%, not less. Process shows technol, effectiveness and can be used in pharmaceutical industry.


- IC ICM A61K031-66
 - ICS A61K031-225; A61K031-785; A61K038-55; A61P031-18
- CC 1-5 (Pharmacology)
- Section cross-reference(s): 63
- IT Amino acids, reactions
 - RL: RCT (Reactant); RACT (Reactant or reagent)
 - (agent for inhibition of reproduction of enveloped viruses, method for its preparing, pharmaceutical composition and method for inhibition of viral infections)
- IT Amino acids, reactions
 - RL: RCT (Reactant); RACT (Reactant or reagent)
 - (salts, potassium and sodium; agent for inhibition of reproduction of enveloped viruses, method for its preparing, pharmaceutical composition and method for inhibition of viral infections)
- IT 99685-96-8DP, Fullerene, homologs
 - RL: ADV (Adverse effect, including toxicity); PAC (Pharmacological
 - activity); PRP (Properties); SPN (Synthetic preparation); THU (Therapeutic
 - use); BIOL (Biological study); PREF (Preparation); USES (Uses)
 (agent for inhibition of reproduction of enveloped viruses, method for its
 - preparing, pharmaceutical composition and method for inhibition of viral infections)
- IT 99685-96-8, Fullerene
 - RL: RCT (Reactant); RACT (Reactant or reagent)
 - (agent for inhibition of reproduction of enveloped viruses, method for its preparing, pharmaceutical composition and method for inhibition of viral infections)
 - 99685-96-8DP, Fullerene, homologs
 - RL: ADV (Adverse effect, including toxicity); PAC (Pharmacological
 - activity); PRP (Properties); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 - (agent for inhibition of reproduction of enveloped viruses, method for its preparing, pharmaceutical composition and method for inhibition of viral infections)
- RN 99685-96-8 ZCAPLUS
- CN [5,6]Fullerene-C60-Ih (CA INDEX NAME)



IT 99685-96-8, Fullerene

- RL: RCT (Reactant); RACT (Reactant or reagent)

- DM 99685-96-8 ZCAPLUS
- CN [5,6]Fullerene-C60-Ih (CA INDEX NAME)



L64 ANSWER 10 OF 24 ZCAPLUS COPYRIGHT 2009 ACS on STN ACCESSION NUMBER: 2004:595195 ZCAPLUS Full-text

DOCUMENT NUMBER: 141:277434

TITLE: Iodo-Controlled Selective Formation of

Pyrrolidino[60]fullerene and Aziridino[60]fullerene from the Reaction between C60 and Amino Acid Esters AUTHOR(S): Zhang, Xiang; Gan, Liangbing; Huang, Shaohua; Shi,

CORPORATE SOURCE:

SOURCE:

Key Laboratory of Bioorganic Chemistry and Molecular, Engineering of the Ministry of Education, College of

Chemistry and Molecular Engineering, Peking University, Beijing, 100871, Peop. Rep. China

Journal of Organic Chemistry (2004), 69(17), 5800-5802

CODEN: JOCEAH; ISSN: 0022-3263

American Chemical Society

PUBLISHER:

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 141:277434

The reaction between glycine Me ester and C60 can be effectively controlled by different iodo-reagents. Addition of DIB ((diacetoxyiodo)benzene) yields the 2,5-bismethoxycarbonyl pyrrolidino[60]fullerene under ultrasonic irradiation; whereas addition of DIB-iodine results in the N-methoxycarbonylmethyl aziridino[60]fullerene under ultrasonic irradiation The reaction of sarcosine Me ester with C60 is similar to that of glycine Me ester under these two conditions. Addition of just iodine to a mixture of sarcosine Me ester and C60 affords the tetra(amino) [60] fullerene epoxide C60(0) ((Me) NCH2COOMe) 4. Possible mechanisms are discussed.

- 27-10 (Heterocyclic Compounds (One Hetero Atom))
- 170501-68-5P 175875-62-4P 760192-22-1P ΙT

RL: SPN (Synthetic preparation); PREP (Preparation)

(iodo-controlled reactions of C60 and amino acid esters under microwave irradiation)


- 760192-22-1P
 - RL: SPN (Synthetic preparation); PREP (Preparation)

(iodo-controlled reactions of C60 and amino acid esters under microwave irradiation)


760192-22-1 ZCAPLUS RN


CN Glycine, N,N',N''',0'''-(9a-oxa-1,9(9a)-homo[5,6]fullerene-C60-Ih6,12,15,18-tetray1)tetrakis[N-methyl-, tetramethyl ester (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 1-B





PAGE 2-B





REFERENCE COUNT: 28 THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L64 ANSWER 11 OF 24 ZCAPLUS COPYRIGHT 2009 ACS on STN ACCESSION NUMBER: 2004:214128 ZCAPLUS Full-text

DOCUMENT NUMBER: 140:423925

TITLE: Different hydroxyl radical scavenging activity of

water-soluble β-alanine C60 adducts
AUTHOR(S): Sun, Tao; Jia, Zhishen; Xu, Zhude
CORPORATE SOURCE: College of Food Science, Shanghai F

CORPORATE SOURCE: College of Food Science, Shanghai Fisheries
University, Shanghai, 200090, Peop. Rep. China
SOURCE: Bioorganic & Medicinal Chemistry Letters (2004),

14(7), 1779-1781

CODEN: BMCLE8; ISSN: 0960-894X

PUBLISHER: Elsevier Science B.V.
DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 140:423925

AB Three C60 derivs. [[C60 (NHCH2CH2CONa]n(H]n], n=1, 5, 9] with different addnl. number of β-alanine were synthesized by the control of relative amount of C60 and β-alanine added. Hydroxyl radical scavenging activity of the adducts was evaluated in a copper-catalyzed Haber-Weiss reaction by chemiluminescence technol. The 50% inhibition concns. (IC50's) of A, B, and C were 147.2 μmol/L, 76.3 μmol/L, and 96.2 μmol/L, resp. The difference should be closely related to the nos. of residual C:C bonds in C60, steric effect and electron-withstanding effect of amino group especially

C 34-2 (Amino Acids, Peptides, and Proteins)

Section cross-reference(s): 22, 25

159510-93-7P 634929-28-5P 635303-26-3P RL: CPS (Chemical process); PEP (Physical, engineering or chemical process); SPN (Synthetic preparation); PREP (Preparation); PROC (Process) (preparation and hydroxyl radical scavenging activity of beta alanine fullerene adducts evaluated in copper-catalyzed Haber-Weiss reaction by chemiluminescence technol.)

IT 159510-93-7P 635303-26-3P

RL: CPS (Chemical process); PEP (Physical, engineering or chemical process); SPN (Synthetic preparation); PREP (Preparation); PROC (Process) (preparation and hydroxyl radical scavenging activity of beta alanine fullerene adducts evaluated in copper-catalyzed Haber-Weiss reaction by chemiluminescence technol.)


RN 159510-93-7 ZCAPLUS

CM 1

CRN 159475-44-2 CMF C87 H105 N9 O18

CO 1 11 0 5 NO 0

CCI IDS



9 T D1_NH_CH2_CH2_CO2H 7


RN 635303-26-3 ZCAPLUS

β-Alanine, N,N',N'',N''',N'''-(decahydro[5,6]fullerene-C60-Ih-CN pentavl)pentakis-, pentasodium salt (9CI) (CA INDEX NAME)

CM 1

CRN 635303-25-2 CMF C75 H85 N5 O10

CCI IDS



5 D1_NH_CH2_CH2_CO2H]

REFERENCE COUNT: 18

THERE ARE 18 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

ACCESSION NUMBER: DOCUMENT NUMBER: TITLE:

INVENTOR(S):

L64 ANSWER 12 OF 24 ZCAPLUS COPYRIGHT 2009 ACS on STN 2003:826902 ZCAPLUS Full-text 140:96298 Method for preparing water-soluble salts of amino acid

derivatives of fullerene Rasnetsov, L. D.; Shvartsman, Ya. Yu.; Lvalina, I. K.; Rasnetsova, B. E.; Karnatsevich, V. L.; Suvorova, O. N.; Kutyreva, V. V.; Shchupak, E. A.; Bazyakina, N.

L.; Makarov, S. G.

PATENT ASSIGNEE(S): Zakrytoe Aktsionernoe Obshchestvo "Desko", Russia

SOURCE: Russ., No pp. given

CODEN: RUXXE7
DOCUMENT TYPE: Patent

LANGUAGE: Russian FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
RU 2213048	C1	20030927	RU 2002-118282	20020708
PRIORITY APPLN. INFO.:			RU 2002-118282	20020708
AB The invention relat	es to	the improved	method for preparing	water- solul

The invention relates to the improved method for preparing water-soluble salts of amino acid derivs. of fullerene that can be used in medicine, pharmacol. and microbiol. Invention describes method for preparing water-soluble salts of amino acid derivs. of fullerene of the general formula HC60NH(CH2)nCOOM wherein C60 is a fullerene ring; M is alkaline metal; n = 1, 3, 5. The method involves interaction of fullerene with amino acid salt in an organic solvent medium at heating and the following isolation of the end product. Interaction reaction is carried out in the presence of low-mol. polyalkylene oxide with mol. mass 150-400 ba. The invention provides reduced process time, and reduced manufacturing cost due to use of inexpensive raw materials.

IC ICM C01B031-02

ICS C07C229-06

CC 49-5 (Industrial Inorganic Chemicals)

T 99685-96-8DP, Fullerene, alkali metal amino acid salt derivs. 645420-16-2P 645420-18-4P 645420-20-8P

645420-22-0P 645420-23-1P 645420-24-2P

RL: IMF (Industrial manufacture); PREP (Preparation)

(method for preparing water-soluble salts of amino acid derivs. of

fullerene)

IT 645420-16-2P 645420-18-4P 645420-20-8P

645420-22-0P 645420-23-1P 645420-24-2P

RL: IMF (Industrial manufacture); PREP (Preparation)

(method for preparing water-soluble salts of amino acid derivs. of fullerene)

RN 645420-16-2 ZCAPLUS

CN Glycine, N-[5,6]fulleren-C60-Ih-1(?H)-yl-, monosodium salt (9CI) (CA INDEX NAME)

CM 1

CRN 645420-15-1

CMF C62 H63 N O2

- *** STRUCTURE DIAGRAM IS NOT AVAILABLE ***
- RN 645420-18-4 ZCAPLUS
- CN Butanoic acid, 4-([5,6]fulleren-C60-Ih-1(?H)-ylamino)-, monosodium salt (9CI) (CA INDEX NAME)

CM 1

CRN 645420-17-3

CMF C64 H67 N O2

- *** STRUCTURE DIAGRAM IS NOT AVAILABLE ***
- RN 645420-20-8 ZCAPLUS
- CN Hexanoic acid, 6-([5,6]fulleren-C60-Ih-1(?H)-ylamino)-, monosodium salt

```
10/559681
    (9CI) (CA INDEX NAME)
    CM 1
    CRN 645420-19-5
     CMF C66 H71 N O2
*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***
RN 645420-22-0 ZCAPLUS
CN Hexanoic acid, 6-([5,6]fulleren-C60-Ih-1(?H)-vlamino)-, monopotassium salt
    (9CI) (CA INDEX NAME)
    CM 1
    CRN 645420-19-5
    CMF C66 H71 N O2
*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***
   645420-23-1 ZCAPLUS
CN Butanoic acid, 4-([5,6]fulleren-C60-Ih-1(?H)-ylamino)-, monopotassium salt
    (9CI) (CA INDEX NAME)
    CM
        1
    CRN 645420-17-3
    CMF C64 H67 N O2
*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***
RN 645420-24-2 ZCAPLUS
CN
    Glycine, N-[5,6]fulleren-C60-Ih-1(?H)-vl-, monopotassium salt (9CI) (CA
    INDEX NAME)
    CM 1
    CRN 645420-15-1
     CMF C62 H63 N O2
*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***
L64 ANSWER 13 OF 24 ZCAPLUS COPYRIGHT 2009 ACS on STN
ACCESSION NUMBER:
                        2003:604154 ZCAPLUS Full-text
DOCUMENT NUMBER:
                        140:42435
TITLE:
                        Effect of functional groups on the activity of
                        water-soluble \beta-alanine C60 derivatives for
                        superoxide anion radical scavenging
AUTHOR(S):
                        Sun, Tao; Xu, Zhu-De; Jia, Zhi-Shen
CORPORATE SOURCE:
                        Dep. Chem., Yuguan Campus, Zhejiang Univ., Hangzhou,
                        310027, Peop. Rep. China
SOURCE:
                        Gaodeng Xuexiao Huaxue Xuebao (2003), 24(7), 1231-1233
                        CODEN: KTHPDM; ISSN: 0251-0790
PUBLISHER:
                        Gaodeng Jiaoyu Chubanshe
DOCUMENT TYPE:
                        Journal
LANGUAGE:
                        Chinese
OTHER SOURCE(S):
                        CASREACT 140:42435
     Three water-soluble \beta-alamine C60 adducts with different addition nos.,
     C60(NHCH2CH2COONa)nHn (n = 1, 5, 9), were synthesized. The products were
     characterized by FTIR, 1H NMR and elemental anal. The antioxidant activity of
     these C60 adducts as the quencher for superoxide anion radical O2- was
     evaluated by chemiluminescence in the system of pyrogallol-luminol. The
```

results indicate that the three C60 derivs. are all excellent quencher for superoxide anion radical O2→4. The concns. of 50% inhibition are 252, 140, 112, µmol/L resp. This high efficiency should be attributed to many factors such as the nos. of remaining double bonds, donor effect of amino group and steric effect. However, the above results also suggest the effect of the adducted β-alanine groups is predominated in this system.

CC 34-2 (Amino Acids, Peptides, and Proteins) Section cross-reference(s): 1, 22

159510-93-7P 634929-28-5P 635303-26-3P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)

(effect of functional groups on activity of water-soluble β -alanine C60 derivs. for superoxide anion radical scavenging)

IT 159510-93-7P 635303-26-3P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)


(effect of functional groups on activity of water-soluble $\beta\text{-alanine}$ C60 derivs. for superoxide anion radical scavenging) 159510-93-7 ZCAPUS

CM 1

RN

CRN 159475-44-2 CMF C87 H105 N9 O18

CCI IDS



RN 635303-26-3 ZCAPLUS


CN β -Alanine, N,N',N'',N''',N'''-(decahydro[5,6]fullerene-C60-Ihpentayl)pentakis-, pentasodium salt (9CI) (CA INDEX NAME)

CM 1

CRN 635303-25-2

CMF C75 H85 N5 O10

CCT IDS



5 D1-NH-CH2-CH2-CO2H

(CA INDEX NAME)

CRN 7601-90-3 CMF C1 H O4

ACCESSION NUMBER:

L64 ANSWER 14 OF 24 ZCAPLUS COPYRIGHT 2009 ACS on STN

DOCUMENT NUMBER: 138:321350 TITLE: Synthesis and characterization of rare-earth fullerene complex AUTHOR(S): Li, Jian-lin; Lin, Yong-sheng; Wu, Zhen-yi; Yang, Sen-gen; Cheng, Da-dian; Zhan, Meng-xiong Dept. of Chem., Xiamen Univ., Xiamen, 361005, Peop. CORPORATE SOURCE: Rep. China Xiamen Daxue Xuebao, Ziran Kexueban (2002), 41(4), SOURCE: 453-455 CODEN: HMHHAF; ISSN: 0438-0479 PUBLISHER: Xiamen Daxue DOCUMENT TYPE: Journal LANGUAGE: Chinese OTHER SOURCE(S): CASREACT 138:321350 AB New rare-earth and buckminsterfullerene[60] complex, C60[La(Gly)2]2(Cl04)6 has been synthesized, and characterized by UV-VIS, IR and elemental anal. The result shows that the rare-earth fullerene complex of n2-form can be synthesized through substituent reaction by using C60 bonding to La(Glv)4Im(ClO4)3. In addition, the structure of the complex was supposed. 29-10 (Organometallic and Organometalloidal Compounds) 511519-43-0P RL: SPN (Synthetic preparation); PREP (Preparation) (synthesis and characterization of lanthanum fullerene glyoxime complex) 511519-43-0P RL: SPN (Synthetic preparation); PREP (Preparation) (synthesis and characterization of lanthanum fullerene glyoxime complex) 511519-43-0 ZCAPLUS RN Lanthanum(2+), [u-[(1,9-n:52,60-n)-[5,6]] fullerene-C60-CN Ih]|tetrakis(qlycinato-κO)di-, diperchlorate, tetraperchlorate (9CI)

2002:630884 ZCAPLUS Full-text

CM 2

CRN 511519-42-9 CMF C68 H16 La2 N4 O8 . 2 C1 O4

CM 3


CRN 511519-41-8

CMF C68 H16 La2 N4 O8 CCI CCS

PAGE 1-A

$$_{\text{H}_2\text{N}-\text{CH}_2-}\mathring{\mathbb{L}}_{-\text{O}^-} \circ - \mathring{\mathbb{L}}_{-\text{CH}_2-\text{NH}_2}$$

PAGE 2-A



PAGE 3-A

$$H_2N-CH_2-\overset{\circ}{\mathbb{U}}_{-0}\overset{-}{-0}\overset{\sum_{a}a}{\longrightarrow} \overset{\circ}{0}-\overset{\circ}{\mathbb{U}}_{-CH_2-NH_2}$$

CM 4

CRN 14797-73-0 CMF C1 O4

·--

CORPORATE SOURCE:

SOURCE:

L64 ANSWER 15 OF 24 ZCAPLUS COPYRIGHT 2009 ACS on STN ACCESSION NUMBER: 2001:902293 ZCAPLUS <u>Full-text</u>

DOCUMENT NUMBER: 136:294749

TITLE: Synthesis and transfection capability of multi-functionalized fullerene polyamine

AUTHOR(S): Isobe, Hiroyuki; Tomita, Naoki; Jinno, Shigeki;

Okayama, Hiroto; Nakamura, Eiichi

Department of Chemistry, Graduate School of Science, The University of Tokyo, Tokyo, 113-0033, Japan

Chemistry Letters (2001), (12), 1214-1215

CODEN: CMLTAG; ISSN: 0366-7022

PUBLISHER: Chemical Society of Japan

DOCUMENT TYPE: Journal LANGUAGE: English

OTHER SOURCE(S): CASREACT 136:294749

AB A new fullerene transfection reagent bearing multiple-functional groups has been synthesized by diastereoselective double cycloaddn. reaction. The highly oxygenated reagent transfers extracellular DNA into mammalian cells with an efficiency comparable to that of a nor-analog.

28-2 (Heterocyclic Compounds (More Than One Hetero Atom))

Section cross-reference(s): 3

IT 407617-27-0P

RL: BCP (Biochemical process); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); PROC (Process)

(preparation and transfection ability of multifunctionalized fullerene polyamine)

IT 407617-27-0P

RL: BCP (Biochemical process); SPN (Synthetic preparation); BIOL

(Biological study); PREP (Preparation); PROC (Process)

(preparation and transfection ability of multifunctionalized fullerene polyamine)

RN 407617-27-0 ZCAPLUS


CN Glycine, N-[3-(dimethylamino)propyl]-N-methyl-, (15,7'R,8'R,95,9'R,135,13'R,145)-8',9'-dihydro-7',13'-dimethoxy-11',11'-dimethyl-5',5''-(ethano|4,5)-endo-[1,3]dioxoloethano)-3'H,3''H-dicyclopenta[1,9:13,14][5,6]fullerene-C60-Ih-3,3''-diyl ester (CA INDEX NAME)


Absolute stereochemistry.


PAGE 1-C















PAGE 4-A



REFERENCE COUNT: 24 THERE ARE 24 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L64 ANSWER 16 OF 24 ZCAPLUS COPYRIGHT 2009 ACS on STN ACCESSION NUMBER: 2001:733394 ZCAPLUS Full-text

DOCUMENT NUMBER: 136:81466

TITLE: Atomic force microscope studies on condensation of plasmid DNA with functionalized fullerenes

AUTHOR(S): Isobe, Hiroyuki; Sugiyama, Sho; Fukui, Ken-ichi;

Iwasawa, Yasuhiro; Nakamura, Eiichi

CORPORATE SOURCE: Department of Chemistry, University of Tokyo, Tokyo, 113-0033, Japan

SOURCE: Angewandte Chemie, International Edition (2001),

40(18), 3364-3367 CODEN: ACIEF5: ISSN: 1433-7851

English

PUBLISHER: Wiley-VCH Verlag GmbH

DOCUMENT TYPE: Journal

LANGUAGE:

AB DNA condensation by a fullerene vector has been imaged at the mol. level by atomic force microscopy (AFM). Anal. of a mixture of plasmid DNA and DNA-binding fullerene with the aid of this carbonaceous vector provided the first information on the mechanism of transfection.

CC 6-2 (General Biochemistry)

T 226420-73-1 226420-75-3

RL: BSU (Biological study, unclassified); BIOL (Biological study) (atomic force microscope studies on condensation of plasmid DNA with functionalized fullerenes)

IT 226420-73-1


RL: BSU (Biological study, unclassified); BIOL (Biological study)
(atomic force microscope studies on condensation of plasmid DNA with
functionalized fullerenes)

RN 226420-73-1 ZCAPLUS

CN Glycine, N-[3-(dimethylamino)propyll-N-methyl-,

1,1'-(5',5''-hexano-3'H,3''H-dicyclopenta[1,9:13,14][5,6]fullerene-C60-Ih-3',3''-diyl) ester (CA INDEX NAME)

$$\begin{array}{c} \text{OLD} & \text{Me} \\ \text{OLD} & \text{In} & \text{CCH}_2 \text{ } \\ \text{3-NMe}_2 \end{array}$$



REFERENCE COUNT:

27 THERE ARE 27 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

ACCESSION NUMBER:

DOCUMENT NUMBER: TITLE:

AUTHOR(S):

CORPORATE SOURCE:

SOURCE:

L64 ANSWER 17 OF 24 ZCAPLUS COPYRIGHT 2009 ACS on STN 2000:894324 ZCAPLUS Full-text

135:71840

Functionalized fullerene as an artificial vector for transfection Nakamura, Eiichi; Isobe, Hiroyuki; Tomita, Naoki;

Sawamura, Masaya; Jinno, Shigeki; Okayama, Hiroto Department of Chemistry, The University of Tokyo, Tokyo, 113-0033, Japan

Angewandte Chemie, International Edition (2000), 39(23), 4254-4257

CODEN: ACIEF5; ISSN: 1433-7851

PUBLISHER: Wiley-VCH Verlag GmbH

DOCUMENT TYPE: Journal LANGUAGE: English

AB The authors synthesized a two-handed fullerene tetramine as well as various related compds. The affinities of these reagents for DNA duplexes were probed. The authors report that the two-handed fullerene tetramine is unique among other fullerenes in its ability to bind to duplex DNA in a reversible manner. Use of the derived fullerene as an artificial vector for transfection is described.

CC 3-2 (Biochemical Genetics)

T 99685-96-8DP, C60 Fullerene, derive. 226420-72-0P 226420-73-1P 226420-74-2P 226420-73-P 226420-76-4P RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)

(functionalized fullerene as artificial vector for transfection)


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)

(functionalized fullerene as artificial vector for transfection)


RN 226420-73-1 ZCAPLUS

CN Glycine, N-[3-(dimethylamino)propyl]-N-methyl-,
1,1'-(5',5''-hexano-3'H,3''H-dicyclopenta[1,9:13,14][5,6]fullerene-C60-Ih3',3''-diyl) ester (CA INDEX NAME)

PAGE 1-A



PAGE 2-A



REFERENCE COUNT: 31 THERE ARE 31 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L64 ANSWER 18 OF 24 ZCAPLUS COPYRIGHT 2009 ACS on STN ACCESSION NUMBER: 2000:634986 ZCAPLUS Full-text

DOCUMENT NUMBER: 133:222456 TITLE: Preparation of amphiphilic fullerenes

INVENTOR(S): Ohishi, Kei; Shinkai, Seiji

PATENT ASSIGNEE(S): Foundation for Scientific Technology Promotion, Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 4 pp.

CODEN: JKXXAF DOCUMENT TYPE: Patent LANGUAGE: Japanese


FAMILY ACC. NUM. COUNT: 1 PATENT INFORMATION:

PATENT NO. KIND DATE APPLICATION NO. DATE -----JP 2000247935 A JP 1999-45704 20000912 19990224 PRIORITY APPLN. INFO.: JP 1999-45704 19990224 OTHER SOURCE(S): MARPAT 133:222456

GI

- AB (FL)Y(O2CXN+Me3BF-)2 [I: FL = fullerene residue; X = (CH2)n, (C2H40)m(CH2)2; Y = Q1, Q2, n = 6-12; m = 2-4]. [60]Fullerene was condensed with 1,2-dibromo-4,5-dimethoxybenzene, demethylated with BBr3, esterified with Br(R2)10COZH, and treated with NMe3 to give I [X = (CH2)10, Y = Q1], which formed a bimol. membrane in H2O.
- IC ICM C07C229-12
- CC 25-29 (Benzene, Its Derivatives, and Condensed Benzenoid Compounds)
- IT 250664-90-5P RL: SPN (Synthetic preparation); PREP (Preparation)
- (preparation of amphiphilic fullerenes)
 T 250664-90-5P
- RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of amphiphilic fullerenes)
- RN 250664-90-5 ZCAPLUS
- CN 1-Undecanaminium, 11,11'-(1',4'-dihydronaphtho[2',3':1,9][5,6]fullerene-C60-Th-6',7'-diyl)bis[N,N,N-trimethyl-11-oxo-, dibromide (9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 2-A

PAGE 3-A

L64 ANSWER 19 OF 24 ZCAPLUS COPYRIGHT 2009 ACS on STN

2000:341607 ZCAPLUS Full-text DOCUMENT NUMBER: 133:120270

ACCESSION NUMBER:

CORPORATE SOURCE:

A novel reaction: [60,70]fullerene reacting with TITLE:

Br-

4,4,5,5-tetramethylimidazoline-2-thione and

α-amino acids as carbene reaction

AUTHOR(S): Xu, Ju-Hua; Li, Yu-Liang; Guo, Zhi-Xin; Li, Feng-Ying; Shi, Zhi-Qiang; Pan, Cai-Yuan; Zhu, Dao-Ben

The Center for Molecular Science, Institute of

Chemistry, Chinese Academy of Sciences, Beijing,

100080, Peop. Rep. China

SOURCE: Journal of Physics and Chemistry of Solids (2000),

61(7), 1081-1088

CODEN: JPCSAW; ISSN: 0022-3697 PUBLISHER:

Elsevier Science Ltd.

DOCUMENT TYPE: Journal

LANGUAGE . English

OTHER SOURCE(S): CASREACT 133:120270

Fullerenes C60 and C70 undergo a [1+2]cycloaddn. reaction with 4,4,5,5tetramethylidazoline-2-thione (I) and DL-valine. C60 was also reacted with I and L-leucine and α -aminoisobutyric acid in order to understand the reaction better and a possible mechanism was put forward based on the results of characterization. In this reaction, reaction temperature was very important due to the reactivity of different α -amino acids and the results also show

that DL-valine was more reactive than the other two α -amino acids.

28-9 (Heterocyclic Compounds (More Than One Hetero Atom))

Amino acids, reactions

RL: RCT (Reactant); RACT (Reactant or reagent)

(reaction of fullerene C60 and C70 with

4.4.5.5-tetramethylimidazoline-2-thione and α-amino acids) 61-90-5, L-Leucine, reactions 62-57-7, 2-Aminoisobutyric acid 516-06-3, Valine 32349-17-0 99685-96-8, [5,6]Fullerene-C60-Ih 115383-22-7, Fullerene C70

ΙT

RL: RCT (Reactant); RACT (Reactant or reagent) (reaction of fullerene C60 and C70 with

4.4.5.5-tetramethylimidazoline-2-thione and α-amino acids)

99685-96-8DP, Fullerene C60, tris adduct with 4, 4, 5, 5-tetramethylimidazolidine-2-thione

RL: SPN (Synthetic preparation); PREP (Preparation)

(reaction of fullerene C60 and C70 with

4.4.5.5-tetramethylimidazoline-2-thione and α-amino acids)

39685-96-8, [5,6]Fullerene-C60-Ih


RL: RCT (Reactant); RACT (Reactant or reagent)

(reaction of fullerene C60 and C70 with

4,4,5,5-tetramethylimidazoline-2-thione and α-amino acids)

99685-96-8 ZCAPLUS RN

CN [5,6]Fullerene-C60-Ih (CA INDEX NAME)




TT 99685-96-8DP, Fullerene C60, tris adduct with 4, 4, 5, 5-tetramethylimidazolidine-2-thione RL: SPN (Synthetic preparation); PREP (Preparation) (reaction of fullerene C60 and C70 with

4,4,5,5-tetramethylimidazoline-2-thione and α -amino acids)

99685-96-8 ZCAPLUS RN

CN [5,6]Fullerene-C60-Ih (CA INDEX NAME)



REFERENCE COUNT: 40 THERE ARE 40 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L64 ANSWER 20 OF 24 ZCAPLUS COPYRIGHT 2009 ACS on STN ACCESSION NUMBER: 2000:161809 ZCAPLUS Full-text

DOCUMENT NUMBER: 132:313992

TITLE: Vesicle Formation and Its Fractal Distribution by Bola-Amphiphilic [60]Fullerene

AUTHOR(S): Sano, Masahito; Oishi, Kei; Ishi, Tsutomu; Shinkai,

Seiji

CORPORATE SOURCE: Chemotransfiguration Project - JST, Kurume, Fukuoka,

839-0861, Japan

SOURCE: Langmuir (2000), 16(8), 3773-3776 CODEN: LANGD5; ISSN: 0743-7463

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal LANGUAGE: English

AB A novel amphiphilic [60]fullerene derivative with two ammonium headgroups is synthesized, and its self-organization characteristics in water in the scale ranging from nanometer to micrometer are reported. At the mol. scale, the bola-amphiphilic [60]fullerene forms spherical vesicles. These vesicles, in turn, are placed within a thin wall producing a foamilke network in the scale-up to a few micrometers. TEM and light scattering measurements demonstrate that the mesoscopic-scale structure is self-similar and fractal with the dimension D = 1.40. The novel aggregation modes result from the hydrophobic interaction produced by the [60]fullerene moieties exposed to water mols. by

C 66-2 (Surface Chemistry and Colloids)
Section cross-reference(s): 3, 22, 73

the disordered alkyl tails.

IT 250664-88-1P 250664-89-2P 250664-90-5P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(vesicle formation and fractal distribution by bola-amphiphilic C60-fullerene derivs.)


IT 250664-90-5P


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(vesicle formation and fractal distribution by bola-amphiphilic C60-fullerene derivs.)

RN 250664-90-5 ZCAPLUS

CN 1-Undecanaminium, 11,11'-(1',4'-dihydronaphtho[2',3':1,9][5,6]fullerene-C60-Ih-6',7'-diyl)bis[N,N,N-trimethyl-11-oxo-, dibromide (9CI) (CA INDEX NAME)





PAGE 2-A

PAGE 3-A

2 Br-

REFERENCE COUNT:

25 THERE ARE 25 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L64 ANSWER 21 OF 24 ZCAPLUS COPYRIGHT 2009 ACS on STN ACCESSION NUMBER: DOCUMENT NUMBER: TITLE:

1999:661948 ZCAPLUS Full-text 131:356589

AUTHOR(S):

Unexpected discovery of a novel organic gel system comprised of [60]fullerene-containing amphiphiles Oishi, Kei; Ishi-I, Tsutomu; Sano, Masahito; Shinkai, Seiii

CORPORATE SOURCE: Chemotransfiguration Project-JST, Fukuoka, 839-0861,

Japan

Chemistry Letters (1999), (10), 1089-1090 SOURCE:

CODEN: CMLTAG; ISSN: 0366-7022

PUBLISHER: Chemical Society of Japan

DOCUMENT TYPE: Journal

LANGUAGE: English

AB A [60]fullerene-containing amphiphile bearing two ammonium groups was synthesized. When the methanol solution was left at room temperature for a few days, it was totally transformed into an organic gel. The transformation process was fully characterized by transmission electron microscopy and X-ray diffraction. This is the first example for the [60]fullerene-containing organic gel.

66-4 (Surface Chemistry and Colloids)

250664-90-5P

RL: PEP (Physical, engineering or chemical process); PRP (Properties); SPN (Synthetic preparation); PREP (Preparation); PROC (Process)

(a novel organic gel system comprised of [60]fullerene-containing amphiphiles

in methanol solution)

250664-90-5P

RL: PEP (Physical, engineering or chemical process); PRP (Properties); SPN (Synthetic preparation); PREP (Preparation); PROC (Process)

(a novel organic gel system comprised of [60]fullerene-containing amphiphiles


in methanol solution)

250664-90-5 ZCAPLUS

CN 1-Undecanaminium, 11,11'-(1',4'-dihydronaphtho[2',3':1,9][5,6]fullerene-C60-Ih-6', 7'-diyl) bis[N, N, N-trimethyl-11-oxo-, dibromide (9CI) (CA INDEX NAME)

PAGE 1-A

PAGE 2-A



PAGE 3-A

2 Br-

REFERENCE COUNT: 10 THERE ARE 10 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L64 ANSWER 22 OF 24 ZCAPLUS COPYRIGHT 2009 ACS on STN ACCESSION NUMBER: 1999:595122 ZCAPLUS Full-text

DOCUMENT NUMBER: 131:237139

TITLE: Fullerene derivatives for potential use in gene

therapy

Nakamura, Eiichi; Sawamura, Masaya; Isobe, Hiroyuki INVENTOR(S):

PATENT ASSIGNEE(S): Fujisawa Pharmaceutical Co., Ltd., Japan

SOURCE: PCT Int. Appl., 37 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Pat.ent. LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PAT	PATENT NO.					KIND		DATE		APPLICATION NO.				DATE				
WO	0 9946235				A1		19990916		WO 1999-JP1146			19990310						
	W:	JP,	US															
	RW:			CH,	CY,	DE,	DK,	ES,	FI,	FR,	GB,	GR,	ΙE,	IT,	LU,	MC,	NL,	
		PT,	SE															
EP	1069107			A1 20010117				EP 1999-907890					19990310					
EP	1069107				B1 20050511													
	R:	AT,	BE,	CH,	DE,	DK,	ES,	FR,	GB,	GR,	IT,	LI,	LU,	NL,	SE,	PT,	ΙE,	FΙ
EP	1420066			A2 20040519				EP 2004-2101					19990310					
EP	1420	066			A3		2005	0105										
	R:	AT,	BE,	CH,	DE,	DK,	ES,	FR,	GB,	GR,	IT,	LI,	LU,	NL,	SE,	PT,	IE,	FΙ
AT	295345				T 20050515					AT 1999-907890				19990310				
US	6765098				B1		20040720			US 2000-622915				20001117				
US	S 20040214218				A1		20041028			US 2004-846646				20040517				
US	7018	599			B2		2006	0328										
PRIORITY	RIORITY APPLN. INFO.:									JP 1	998-	5861	4		A 1	9980	310	
										EP 1	999-	9078	90		A3 1	9990	310	

WO 1999-JP1146 W 19990310 US 2000-622915 A3 20001117

OTHER SOURCE(S): MARPAT 131:237139

AB This document discloses a novel means for DNA compaction. Fullerene derivs. having 1 to 4 nitrogen-containing hydrophilic side chains or salts thereof are to be used in the above means. A fullerene derivative was prepared and tested using calf thymus DNA and ethidium bromide.

IC ICM C07C229-16

ICS C12N015-00; A61K048-00

CC 78-1 (Inorganic Chemicals and Reactions)

Section cross-reference(s): 1, 3 IT 226420-73-1P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses) (fullerene derivs, for potential use in dene therapy)

IT 226420-73-1P


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses) (fullerene derive. for potential use in gene therapy)

RN 226420-73-1 ZCAPLUS

CN Glycine, N-[3-(dimethylamino)propyl]-N-methyl-, 1,1'-(5',5''-hexano-3'H,3''H-dicyclopenta[1,9:13,14][5,6]fullerene-C60-Ih-3',3''-diyl) ester (CA INDEX NAME)

PAGE 1-A

PAGE 2-A



REFERENCE COUNT:

THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD, ALL CITATIONS AVAILABLE IN THE RE FORMAT

ACCESSION NUMBER:

DOCUMENT NUMBER:

TITLE:

AUTHOR(S):

CORPORATE SOURCE:

SOURCE:

PUBLISHER: DOCUMENT TYPE: LANGUAGE:

GI

L64 ANSWER 23 OF 24 ZCAPLUS COPYRIGHT 2009 ACS on STN

1999:285374 ZCAPLUS Full-text

131:18823 Synthesis and functions of a novel DNA binding


fullerene Isobe, Hiroyuki; Sawamura, Masaya; Sugiyama, Sho; Fukui, Ken-ich; Iwasawa, Yasuhiro; Nakamura, Eiichi Department of Chemistry, The University of Tokyo,

Japan Tennen Yuki Kagobutsu Toronkai Koen Yoshishu (1998),

40th, 157-161


CODEN: TYKYDS Nippon Kagakkai Journal

Japanese




- AB Much interest in the biol. activities of fullerenes have been drawn since their first discovery. As one of the first findings of biol. activities, we have already reported the synthesis of water-miscible fullerene carboxylic acid and its biol. activities. Thus, we found that fullerenes show the photoinduced DNA cleaving activity, cytotoxicity and enzyme inhibition. In this study, we further investigated their biol. activities, including distribution anal. of dosed radio-active fullerene and interactions with DNAs. We have synthesized radio-active water-miscible fullerene carboxylic acid by [3 + 2] cycloaddn, of trimethylene methane. When administered orally to rats, fullerene was not efficiently absorbed and was excreted primarily in the feces. When injected i.v., it was distributed rapidly to various tissues, and most of the material was retained in the body after one week. We have expanded fullerene's photo-induced biol. activities, targeting DNAs. Thus, we have designed and synthesized a novel fullerene-oligonucleotide conjugate, and examined its DNA nicking ability. When this conjugate was incubated and irradiated with oligonucleotides with target seguence, site specific nicking of the oligonucleotide was achieved. We further continued investigating the interactions of fullerene with DNAs and found that fullerene-polyamine [I; R = COCH2NMe(CH2)3NMe2| conjugate tightly binds to DNA in a non-specific manner, and this new fullerene gave first example of photo-induced covalent bond formation of fullerene with DNA base.
- CC 25-29 (Benzene, Its Derivatives, and Condensed Benzenoid Compounds) Section cross-reference(s): 6
- IT 226420-72-0P 226420-73-1P 226420-74-2P 226420-75-3P
 Rl: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 - (synthesis and functions of novel DNA binding fullerene)
- IT 226420-73-1P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 - (synthesis and functions of novel DNA binding fullerene)
- RN 226420-73-1 ZCAPLUS
- CN Glycine, N-[3-(dimethylamino)propyl]-N-methyl-, 1,1'-(5',5''-hexano-3'',3''H-dicyclopenta[1,9:13,14][5,6]fullerene-C60-Ih-3',3''-divl) ester (CA INDEX NAME)

PAGE 1-A







L64 ANSWER 24 OF 24 ZCAPLUS COPYRIGHT 2009 ACS on STN ACCESSION NUMBER: 1995:20078 ZCAPLUS Full-text

DOCUMENT NUMBER:

122:10488

ORIGINAL REFERENCE NO.: 122:2333a,2336a

TITLE:

SOURCE:

Water-soluble fullerene derivatives. Synthesis and

characterization of \$\beta\$-alanine C60 adducts

AUTHOR(S): Gan, Liang Bing; Luo, Chu Ping; Xu, Lian Bin; Zhou, De

Jing; Huang, chun Hui; Zhao, Shan Kai

CORPORATE SOURCE: Department Chemistry, Peking University, Beijing,

> 100871, Peop. Rep. China Chinese Chemical Letters (1994), 5(4), 275-8

CODEN: CCLEE7: ISSN: 1001-8417

Journal

DOCUMENT TYPE: LANGUAGE: English

OTHER SOURCE(S): CASREACT 122:10488

AB β -Alanine sodium salt reacts with C60 to give a water-soluble derivative C60(NHCH2CH2CO2Na)x(H)x (I). Acidification of I yields derivative C60(NHCH2CH2CO2H)x(H)x. Elemental anal. suggest x is equal to 9. 1H and 13C NMR spectra are consistent with the addition of the amino acid through its amino group. These amino acid derivs, are air-stable and may be used as precursors for further functionalization.

CC 34-2 (Amino Acids, Peptides, and Proteins)

Section cross-reference(s): 24, 78

159475-45-3P

RL: SPN (Synthetic preparation); PREP (Preparation)

(preparation and acidification of)

159510-93-7P

RL: SPN (Synthetic preparation); PREP (Preparation)

(preparation of)

159475-45-3P

RL: SPN (Synthetic preparation); PREP (Preparation)


(preparation and acidification of)

RN 159475-45-3 ZCAPLUS

β-Alanine, N,N',N'',N''',N'''',N''''',N''''',N'''''-(octahydro[5,6]fullerene-C60-Ih-nonayl)nonakis- (9CI) (CA INDEX NAME) CM 1

CRN 159475-44-2 CMF C87 H105 N9 O18

CCT TDS



IT 159510-93-7P

RL: SPN (Synthetic preparation); PREP (Preparation)

(preparation of)


RN 159510-93-7 ZCAPLUS

β-Alanine, N,N',N'',N''',N'''',N'''',N''''',N'''''-CN (octahydro[5,6]fullerene-C60-Ih-nonayl)nonakis-, nonasodium salt (9CI) (CA INDEX NAME)

CM 1

CRN 159475-44-2 CMF C87 H105 N9 O18

CCT IDS



9
$$\square$$
 D1_NH_CH2_CH2_CO2H \square

=> d his full (FILE 'HOME' ENTERED AT 15:13:50 ON 02 JAN 2009) FILE 'ZCAPLUS' ENTERED AT 15:14:11 ON 02 JAN 2009 E US2005-559681 /APPS 1 SEA SPE=ON ABB=ON PLU=ON US2005-559681 /AP D SCA FILE 'REGISTRY' ENTERED AT 15:16:32 ON 02 JAN 2009 1 SEA SPE=ON ABB=ON PLU=ON 99685-96-8 D SCA FILE 'ZCAPLUS' ENTERED AT 15:16:57 ON 02 JAN 2009 T. 3 2246 SEA SPE=ON ABB=ON PLU=ON L2/D L4 977 SEA SPE=ON ABB=ON PLU=ON L3 (L) PREP/RL 207299 SEA SPE=ON ABB=ON PLU=ON AMINO ACID?/CW L5 15098 SEA SPE=ON ABB=ON PLU=ON L5 (L) RACT/RL L6 L7 4881 SEA SPE=ON ABB=ON PLU=ON L2 (L) RACT/RL L8 30 SEA SPE=ON ABB=ON PLU=ON L6 AND L7 2 SEA SPE=ON ABB=ON PLU=ON L4 AND L8 L9 D SCA T.10 28 SEA SPE=ON ABB=ON PLU=ON L8 NOT L9 118009 SEA SPE=ON ABB=ON PLU=ON ?DIAMINO?/BI OR ?TRIAMINO?/BI OR L11 DI AMINO?/BI OR ?TRI AMINO?/BI 1 SEA SPE=ON ABB=ON PLU=ON L8 AND L11 D SCA D KWIC D BIB 6 SEA SPE=ON ABB=ON PLU=ON L8 AND P/DT T.13 24 SEA SPE=ON ABB=ON PLU=ON L8 NOT L13 L14 L*** DEL 1 S L13 AND PY<2004 L15 13 SEA SPE=ON ABB=ON PLU=ON L14 AND PY<2004 FILE 'REGISTRY' ENTERED AT 15:33:42 ON 02 JAN 2009 L16 STRUCTURE UPLOADED L17 5 SEA SSS SAM L16 D SCA D STAT OUE STRUCTURE UPLOADED L18 L19 STRUCTURE UPLOADED L20 5 SEA SSS SAM L18 AND L19 D SCA D STAT QUE L21 260 SEA SSS FUL L18 AND L19 SAVE TEMP JEA681L18L19/A L21 L22 37 SEA SPE-ON ABB-ON PLU-ON L21 AND IDS/CI 13 SEA SPE=ON ABB=ON PLU=ON L22 AND C60/CNS 1.23 L24 24 SEA SPE=ON ABB=ON PLU=ON L22 AND C82/CNS L25 37 SEA SPE=ON ABB=ON PLU=ON L23 OR L24 D SCA L24 FILE 'ZCAPLUS' ENTERED AT 15:46:53 ON 02 JAN 2009 6 SEA SPE=ON ABB=ON PLU=ON L23 L26 D IBIB HITSTR 6

FILE 'REGISTRY' ENTERED AT 15:48:01 ON 02 JAN 2009
223 SEA SPE=ON ABB=ON PLU=ON L21 NOT L22

```
10/559681
1.28
            39 SEA SPE=ON ABB=ON PLU=ON L27 AND NC>1
               D SCA
1.29
               STRUCTURE UPLOADED
L30
             6 SEA SUB=L21 SSS SAM L29
               D SCA
L31
               STRUCTURE UPLOADED
L32
             0 SEA SUB=L21 SSS SAM L31
L33
             0 SEA SUB=L21 SSS FUL L31
              D STAT OUE L30
L34
            70 SEA SUB=L21 SSS FUL L29
L35
              STRUCTURE UPLOADED
L36
               STRUCTURE UPLOADED
            0 SEA SUB=L21 SSS SAM L35 OR L36
L37
1.38
            0 SEA SUB=L21 SSS FUL L35 OR L36
             STRUCTURE UPLOADED
L39
L40
              STRUCTURE UPLOADED
            2 SEA SUB=L21 SSS SAM L39 OR L40
L41
               D SCA
L42
            15 SEA SUB=L21 SSS FUL L39 OR L40
               D SCA
    FILE 'ZCAPLUS' ENTERED AT 16:06:54 ON 02 JAN 2009
             15 SEA SPE=ON ABB=ON PLU=ON L42
               SET NOTICE OFF DISPLAY
               SET NOTICE OFF SEARCH
           22 SEA SPE=ON ABB=ON PLU=ON RASNETSOV L?/AU
55 SEA SPE=ON ABB=ON PLU=ON SHVARTSMAN I?/AU
T. 44
L45
L46
           13 SEA SPE=ON ABB=ON PLU=ON LYALINA I?/AU
           19 SEA SPE=ON ABB=ON PLU=ON RASNETSOVA B?/AU
L47
            O SEA SPE=ON ABB=ON PLU=ON L44 AND L45 AND L46 AND L47
L48
           15 SEA SPE=ON ABB=ON PLU=ON L44 AND (L45 OR L46 OR L47)
L49
L50
            1 SEA SPE=ON ABB=ON PLU=ON L45 AND (L46 OR L47)
            2 SEA SPE=ON ABB=ON PLU=ON L46 AND L47
15 SEA SPE=ON ABB=ON PLU=ON L49 OR L50 OR L51
L51
L52
               SET NOTICE LOGIN DISPLAY
               SET NOTICE LOGIN SEARCH
   FILE 'MEDLINE, EMBASE, BIOSIS, WPIX' ENTERED AT 16:09:13 ON 02 JAN 2009
L53
            14 SEA SPE=ON ABB=ON PLU=ON L52
    FILE 'ZCAPLUS' ENTERED AT 16:09:35 ON 02 JAN 2009
L54
         38813 SEA SPE=ON ABB=ON PLU=ON ?FULLEREN?/BI
L55
            11 SEA SPE=ON ABB=ON PLU=ON (L44 OR L45 OR L46 OR L47) AND L54
            18 SEA SPE=ON ABB=ON PLU=ON L55 OR L52
L57
             2 SEA SPE=ON ABB=ON PLU=ON L5 AND (L52 OR L55)
    FILE 'MEDLINE, EMBASE, BIOSIS, WPIX' ENTERED AT 16:10:59 ON 02 JAN 2009
L58
             9 SEA SPE-ON ABB-ON PLU-ON (L44 OR L45 OR L46 OR L47) AND
               ?FULLEREN?
```

SEA SPEON ABB-ON PLU-ON (L53 OR L58) AND AMINO ACID?

FILE 'ZCAPLUS' ENTERED AT 16:11:56 ON 02 JAN 2009

D STAT QUE L52

D STAT QUE L55

D STAT QUE L57 L60 18 SEA SPE=ON ABB=ON PLU=ON L52 OR L55 OR L52

L59

FILE 'MEDLINE, EMBASE, BIOSIS, WPIX' ENTERED AT 16:12:21 ON 02 JAN 2009

D STAT QUE L53

D STAT QUE L58

D STAT QUE L59

L61 14 SEA SPE=ON ABB=ON PLU=ON L53 OR L58 OR L59

FILE 'ZCAPLUS, WPIX' ENTERED AT 16:12:55 ON 02 JAN 2009
19 DUP REM L60 L61 (13 DUPLICATES REMOVED)

ANSWERS '1-18' FROM FILE ZCAPLUS ANSWER '19' FROM FILE WPIX

D IBIB ABS HITIND L62 1-18

IBIB ABS HITIND L62 1-

D IALL HIT L62 19

FILE 'REGISTRY' ENTERED AT 16:13:36 ON 02 JAN 2009

FILE 'ZCAPLUS' ENTERED AT 16:13:40 ON 02 JAN 2009

D STAT QUE L26 D STAT OUE L43

L63 21 SEA SPE=ON ABB=ON PLU=ON L26 OR L43

D STAT QUE L9 D STAT OUE L12

L64 24 SEA SPE=ON ABB=ON PLU=ON L26 OR L43 OR L9 OR L12
D IBIB ABS HITIND HITSTR L64 1-24

FILE HOME

FILE ZCAPLUS

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS is strictly prohibited.

FILE COVERS 1907 - 2 Jan 2009 VOL 150 ISS 2 FILE LAST UPDATED: 1 Jan 2009 (20090101/ED)

ZCAplus now includes complete International Patent Classification (IPC) reclassification data for the third quarter of 2008.

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file contains CAS Registry Numbers for easy and accurate substance identification.

FILE REGISTRY

Property values tagged with IC are from the ${\tt ZIC/VINITI}$ data file provided by InfoChem.

STRUCTURE FILE UPDATES: 1 JAN 2009 HIGHEST RN 1092443-48-5 DICTIONARY FILE UPDATES: 1 JAN 2009 HIGHEST RN 1092443-48-5

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH July 5, 2008.

Please note that search-term pricing does apply when

conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

http://www.cas.org/support/stngen/stndoc/properties.html

FILE MEDLINE

FILE LAST UPDATED: 1 Jan 2009 (20090101/UP). FILE COVERS 1949 TO DATE.

MEDLINE and LMEDLINE have been updated with the 2009 Medical Subject Headings (MeSH) vocabulary and tree numbers from the U.S. National Libra of Medicine (NLM). Additional information is available at

http://www.nlm.nih.gov/pubs/techbull/nd08/nd08_medline_data_changes_2009.

In preparation for the annual MEDLINE reload, NLM suspends delivery of regular updates (completed records), but continues to send "in-process" records. STN will resume regular MEDLINE updates the week of Dec 29, 200

This file contains CAS Registry Numbers for easy and accurate substance identification.

See HELP RANGE before carrying out any RANGE search.

MEDLINE Accession Numbers (ANs) for records from 1950-1977 have been converted from 8 to 10 digits. Searches using an 8 or 10 digit AN will retrieve the same record. The 10-digit ANs can be expanded, searched, and displayed in all records from 1949 to the present.

FILE EMBASE

FILE COVERS 1974 TO 31 Dec 2008 (20081231/ED)

EMBASE was reloaded on March 30, 2008.

EMBASE is now updated daily. SDI frequency remains weekly (default) and biweekly.

This file contains CAS Registry Numbers for easy and accurate substance identification.

Beginning January 2008, Elsevier will no longer provide EMTREE codes as part of the EMTREE thesaurus in EMBASE. Please update your current-awareness alerts (SDIs) if they contain EMTREE codes.

For further assistance, please contact your local helpdesk.

FILE BIOSIS

FILE COVERS 1926 TO DATE.

CAS REGISTRY NUMBERS AND CHEMICAL NAMES (CNs) PRESENT FROM JANUARY 1926 TO DATE.

RECORDS LAST ADDED: 31 December 2008 (20081231/ED)

BIOSIS has been augmented with 1.8 million archival records from 1926 through 1968. These records have been re-indexed to match current BIOSIS indexing.

FILE WPIX

FILE LAST UPDATED: 22 DEC 2008 <20081222/UP> MOST RECENT UPDATE: 200882 <200882/DW>

DERWENT WORLD PATENTS INDEX SUBSCRIBER FILE, COVERS 1963 TO DATE

>>> Now containing more than 1.2 million chemical structures in DCR <<<

>>> IPC Reform backfile reclassifications have been loaded to end of September 2008. No update date (UP) has been created for the reclassified documents, but they can be identified by 20060101/UPIC, and 20061231/UPIC, 20070601/UPIC, 20071001/UPIC, 20071130/UPIC, 20080401/UPIC, 20080701/UPIC and 20081001/UPIC. ECLA reclassifications to mid August and US national classification

mid September 2008 have also been loaded. Update dates 20080401, 20080701 and 20081001/UPEC and /UPNC have been assigned to these. <<

FOR A COPY OF THE DERWENT WORLD PATENTS INDEX STN USER GUIDE, PLEASE VISIT:

http://www.stn-international.de/training_center/patents/stn_quide.pdf

FOR DETAILS OF THE PATENTS COVERED IN CURRENT UPDATES, SEE http://scientific.thomsonreuters.com/support/patents/coverage/latestupdate

EXPLORE DERWENT WORLD PATENTS INDEX IN STN ANAVIST, VERSION 2.0: http://www.stn-international.com/DWPIAnaVist2 0608.html


>>> HELP for European Patent Classifications see HELP ECLA, HELP ICO <<<



chain nodes : 1 2 3 4 5 chain bonds : 1-2 2-3 3-4 3-5 exact/norm bonds : 3-4 3-5 exact bonds : 1-2 2-3

Connectivity: 2:2 E exact RC ring/chain Match level : 1:CLASS 2:CLASS 3:CLASS 4:CLASS 5:CLASS Generic attributes : 2: Saturation : Saturated

Uploading L19.str



ring nodes :

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20

ring bonds :

1-2 1-6 1-19 2-3 2-9 3-4 3-12 4-5 5-6 6-7 7-8 8-20 9-10 9-13 10-11 10-16 11-12 13-14 13-19 14-15 14-18 15-16 17-18 17-20 19-20


exact/norm bonds :

 $1-2 \quad 1-6 \quad 1-19 \quad 2-3 \quad 2-9 \quad 3-4 \quad 3-12 \quad 4-5 \quad 5-6 \quad 6-7 \quad 7-8 \quad 8-20 \quad 9-10 \quad 9-13 \quad 10-11$

10-16 11-12 13-14 13-19 14-15 14-18 15-16 17-18 17-20 19-20

Match level :

1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom 11:Atom 12:Atom 13:Atom 14:Atom 15:Atom 16:Atom 17:Atom 18:Atom 19:Atom 20:Atom



1 2 3 4 5 7 8 9 10 11 12 chain bonds :

chain nodes :

1-2 1-12 2-3 3-4 3-5 7-8 7-12 8-9 9-10 9-11

exact/norm bonds :


1-12 3-4 3-5 7-8 7-12 8-9 9-10 9-11

exact bonds :

1-2 2-3

```
Connectivity:
2:2 E exact RC ring/chain
Match level:
1:CLASS 2:CLASS 3:CLASS 4:CLASS 5:CLASS 7:CLASS 8:CLASS 9:CLASS 10:CLASS
11:CLASS 12:Atom
Generic attributes:
2:
Saturation: Saturated
```

Element Count : Node 12: Limited C,C55



```
chain nodes:
1 2 3 4 5 7 8 9 10 11 12
chain bonds:
1-2 2-3 3-4 3-5 5-12 7-8 8-9 9-10 9-11 11-12
exact/norm bonds:
3-4 3-5 5-12 7-8 8-9 9-10 9-11 11-12
exact bonds:
1-2 2-3
```

```
Connectivity:
2:2 E exact RC ring/chain
Match level:
1:CLASS 2:CLASS 3:CLASS 4:CLASS 5:CLASS 7:CLASS 8:CLASS 9:CLASS 10:CLASS
11:CLASS 12:Atom
Generic attributes:
2:
Saturation : Saturated
```

Element Count : Node 12: Limited C,C55